

Migraine linked to increased stroke risk in women

November 15 2016



Credit: Sasha Wolff/Wikipedia

Women who have migraines may have an increased risk of stroke, according to a preliminary study presented at the American Heart

Association's Scientific Sessions 2016.

Researchers studied 917 women who were being evaluated for heart disease and found those that had a history of migraines (224 or nearly 25 percent) were at higher risk of a future cardiovascular event - notably [stroke](#).

They also found:

- Compared to those who did not report a history of migraines, women with a history of the headaches had an 83 percent higher risk of a [cardiovascular event](#), including stroke or heart attack, during an average six-year follow-up.
- Women with history of migraine were 2.33 times more likely to suffer a stroke during the study than women who didn't report migraines.

More information: Cecil A. Rambarat, M.D., University of Florida, Gainesville, Florida. (Actual presentation time for T2125 is 1:30 p.m. CT/2:30 p.m. ET, Tuesday, Nov. 15, 2016.)

Provided by American Heart Association

Citation: Migraine linked to increased stroke risk in women (2016, November 15) retrieved 26 December 2022 from <https://medicalxpress.com/news/2016-11-migraine-linked-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.