

Interpersonal violence increases the risk of repeated suicide attempts

17 November 2016

A study conducted by researchers from the Karolinska Institute shows that patients with an experience of violence have an increased risk to carry out repeated suicide attempts.

In a multicenter study conducted at the emergency hospitals in Stockholm, Gothenburg and Umeå, patients who have survived a suicide attempt have been interviewed. The interviews include their experiences of using [violence](#) and being exposed to violence. Of the 355 patients included in the study 78 repeated a [suicide attempt](#) during the follow-up period of six months. The risk of repeating an attempt proved to be associated with their reported experiences of violence. A specific finding among the women in the study was that exposure to violence as an adult was associated with repeated violent suicide attempts.

"Our results support the hypothesis that experiences of violence could be associated with the development of [suicidal behavior](#). Better identification of patients' experiences of violence may enhance suicide risk assessments," says researcher Axel Haglund at the Karolinska Institute.

More information: Axel Haglund et al. Interpersonal violence and the prediction of short-term risk of repeat suicide attempt, *Scientific Reports* (2016). DOI: [10.1038/srep36892](https://doi.org/10.1038/srep36892)

Provided by Karolinska Institutet

APA citation: Interpersonal violence increases the risk of repeated suicide attempts (2016, November 17) retrieved 1 May 2021 from <https://medicalxpress.com/news/2016-11-interpersonal-violence-suicide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.