

# Insomnia prevalent in patients with asthma

8 December 2016

A team of researchers from the University of Pittsburgh has found that insomnia is highly prevalent in adults with asthma and is also associated with worse asthma control, depression and anxiety symptoms and other quality of life and health issues. The study results are published in the current issue of the journal *CHEST*.

Patients with [asthma](#) commonly report difficulty with sleep; however, the prevalence of [insomnia](#) and its relationship with asthma burden and quality of life is unknown. Asthma is a chronic respiratory condition in the lungs in which the airways become swollen or inflamed, causing difficulty in breathing from spasms in the muscles surrounding the airways as they try to keep the passageways open. The impact of insomnia on [asthma control](#) and asthma-related healthcare utilization has not previously been examined.

The research reports that:

- Clinically significant insomnia was present in 37 percent of the participants with asthma.
- Those with insomnia had a higher BMI, worse lung function and lower annual household income than those without insomnia.
- Despite reporting no nighttime asthma symptoms that disturbed their sleep, almost 25 percent of participants met criteria for clinically significant insomnia.
- Compared with those without insomnia, the study found participants with insomnia reported more frequent asthma-related healthcare use in the past 12 months.
- Participants with insomnia had worse asthma control and asthma-specific quality of life and higher levels of depression and [anxiety symptoms](#). These results suggest that adults with asthma who have insomnia disorder may be at increased risk for [adverse outcomes](#).

The research shows that there is a significant

impact of insomnia on asthma disease burden and well-being and states that evaluation and treatment of insomnia should be considered among patients with asthma. "Our results show that poor sleep may not be solely due to nighttime awakenings due to asthma symptoms but may represent comorbid insomnia," said lead author Faith Luyster, PhD, "and that comorbid insomnia can significantly impact asthma outcomes including quality of life and healthcare utilization."

While it was determined that insomnia is highly prevalent in those with asthma and is associated with adverse outcomes, further studies are needed to better understand the relationship between insomnia and asthma control. Prospective and interventional studies, such as implementing cognitive-behavioral treatment for insomnia, are recommended moving forward.

The complete report on this research can be found in the journal *CHEST*.

Provided by American College of Chest Physicians

APA citation: Insomnia prevalent in patients with asthma (2016, December 8) retrieved 5 May 2021 from <https://medicalxpress.com/news/2016-12-insomnia-prevalent-patients-asthma.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*