

Self-care tools cut depression in AMD, diabetic retinopathy

14 December 2016



outcomes, including anxiety, life space, and self-efficacy.

"Self-care tools plus telephone coaching led to a modest improvement in [depressive symptoms](#) in patients with age-related eye disease," the authors write. "Additional research on how to maximize their effect is necessary."

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—Self-care tools can reduce depressive symptoms in patients with age-related macular degeneration (AMD) or diabetic retinopathy (DR), according to a study published online Dec. 7 in *Clinical & Experimental Ophthalmology*.

Hortence Kamga, from Maisonneuve-Rosemont Hospital in Montreal, and colleagues conducted a single-blind randomized controlled clinical trial involving 80 participants with late stage AMD or DR, at least mild depressive symptoms, and [visual acuity](#) better than 20/200. Participants were randomized in a 1:1 ratio to the intervention, which consisted of large-print written and audio tools incorporating cognitive-behavioral principles plus three 10-minute telephone calls, or delayed intervention/usual care.

The researchers found that the intervention reduced depressive symptoms by 2.1 points more than usual care, after adjustment for baseline imbalances in visual acuity ($P = 0.040$). The intervention was not associated with secondary

APA citation: Self-care tools cut depression in AMD, diabetic retinopathy (2016, December 14) retrieved 7 September 2022 from https://medicalxpress.com/news/2016-12-self-care-tools-depression-amd-diabetic_1.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.