


Case volume impacts re-op rates after penile prosthesis Sx

20 December 2016



implants placed by lower-volume implanters.

Compared with patients <u>treated</u> by <u>surgeons</u> in the highest quartile of annual case volume (more than 31 cases per year), those treated by surgeons in the lowest, second, and third annual case volume quartiles (zero to two, three to seven, and eight to 31 cases per year, respectively) were significantly more likely to require reoperation for inflatable penile prosthesis infection (hazard ratios, 2.5, 2.4, and 2.1, respectively).

"Patients treated by higher-volume implanters are less likely to require reoperation after inflatable penile prosthesis insertion than those treated by lower-volume surgeons," the authors write.

One author disclosed financial ties to Coloplast.

More information: Full Text

(HealthDay)—Surgeon volume is associated with the likelihood of reoperation after inflatable penile prosthesis surgery, with the trend driven by reoperations for infection, according to a study published in the January issue of *The Journal of Urology*.

Ifeanyi C. Onyeji, from the Columbia University Medical Center in New York City, and colleagues queried the New York Statewide Planning and Research Cooperative System database from 1995 to 2014 for inflatable penile prosthesis cases. They examined the impact of surgeon annual case volume on inflatable penile prosthesis reoperation rates among 14,969 men followed for a median of 95.1 months from the time of implant.

The researchers found that the rate of overall reoperation was 6.4 percent, and the rates of reoperation from <u>infection</u> and noninfectious complications were 2.5 and 3.9 percent, respectively. Reoperation for infection, but not for noninfectious complications, was more likely for

Copyright © 2016 HealthDay. All rights reserved.



APA citation: Case volume impacts re-op rates after penile prosthesis Sx (2016, December 20) retrieved 5 May 2021 from <u>https://medicalxpress.com/news/2016-12-case-volume-impacts-re-op-penile.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.