

# Higher AlphaB-crystallin levels linked to diabetic retinopathy

21 December 2016



with PDR and present a crucial association between alphaB-crystallin and VEGF with angiogenic activity in PDR," the authors write.

**More information:** [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—Vitreous levels of alphaB-crystallin are significantly higher in eyes of patients with proliferative diabetic retinopathy (PDR) compared to the eyes of controls without diabetes, according to a study published online Dec. 7 in *Clinical & Experimental Ophthalmology*.

Wei Chen, Ph.D., from the Affiliated Hospital of Nantong University in China, and colleagues assessed vitreous samples collected before vitrectomy from 46 eyes of 46 consecutive patients with PDR and 19 patients without [diabetes mellitus](#) who had vitrectomy for idiopathic macular hole. Enzyme-linked immunosorbent assay was used to measure concentrations of alphaB-crystallin and vascular endothelial growth factor (VEGF).

The researchers found that the vitreous level of alphaB-crystallin was significantly higher in patients with PDR versus controls (P

"These results suggest a significant increase of alphaB-crystallin in the vitreous fluid of patients

APA citation: Higher AlphaB-crystallin levels linked to diabetic retinopathy (2016, December 21) retrieved 27 April 2021 from <https://medicalxpress.com/news/2016-12-higher-alphab-crystallin-linked-diabetic-retinopathy.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*