


Opicapone as levodopa adjunct cuts motor fluctuations in Parkinson's disease

28 December 2016



the 25- and 50-mg/day opicapone groups, respectively. For the 50-mg/day opicapone group, the adjusted [treatment](#) difference versus placebo was significant (± 54.3 minutes; $P = 0.008$); the difference was not significant for the 25-mg group (± 37.2 minutes; $P = 0.11$). Throughout the open-label phase, the off-time reduction was sustained (± 126.3 minutes at one-year end point).

"Treatment with a 50-mg once-daily dose of opicapone was associated with a significant reduction in mean daily off-time in levodopa-treated patients with PD and motor fluctuations, and this effect is maintained for at least one year," the authors write.

Several authors disclosed financial ties to pharmaceutical companies, including BIAL, which manufactures opicapone and funded the study.

(HealthDay)—For patients with Parkinson's disease (PD) receiving levodopa therapy and experiencing end-of-dose motor fluctuations, treatment with 50-mg/day opicapone is associated with a reduction in mean daily off-time, according to a study published online Dec. 27 in *JAMA Neurology*.

Andrew J. Lees, M.D., from University College London, and colleagues enrolled 427 [patients](#) with PD receiving [levodopa therapy](#) who experienced signs of end-of-dose deterioration into a phase 3 trial. Participants were randomized to a 14- to 15-week [placebo](#)-controlled trial (129 received 25-mg/day opicapone, 154 received 50-mg/day opicapone, and 144 received placebo). Of the 376 patients who completed the double-blind phase, 286 completed a one-year open-label phase in which all patients received opicapone.

The researchers found that the least squares mean change in off-time was ± 64.5 minutes for the placebo group, and ± 101.7 and ± 118.8 minutes for

More information: [Full Text \(subscription or payment may be required\)](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

APA citation: Opicapone as levodopa adjunct cuts motor fluctuations in Parkinson's disease (2016, December 28) retrieved 5 October 2022 from <https://medicalxpress.com/news/2016-12-opicapone-levodopa-adjunct-motor-fluctuations.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.