


A new index for the diagnosis of non-alcoholic fatty liver disease

24 January 2017



The Fatty Liver Index (FLI) consists of the parameters age, waist circumference, and triglyceride levels (TG) measured in the blood in a fasting state, and gamma-glutamyl transpeptidase (GGT). Credit: © DZD/IDM

Non-alcoholic fatty liver disease has become a global epidemic. Researchers worldwide aim to understand the causes and consequences of fatty liver disease, but also to diagnose fatty liver disease at an early stage. Under the direction of Professor Norbert Stefan, researchers of the Department of Internal Medicine IV of Tübingen University Hospital and their collaborators generated a new index from clinical data that can predict the presence of fatty liver disease with high accuracy.

Approximately every third adult in the industrialized countries has a morbidly fatty liver. This not only increases the risk of advanced liver diseases such as [liver cirrhosis](#) and [liver cancer](#), but also type 2 diabetes and cardiovascular diseases. To be able to intervene early with preventive and therapeutic measures, the fatty liver must be detected in time. For this purpose, ultrasound examination of the

liver and the determination of the liver values are suitable, but in most cases, they can only diagnose the fatty liver in an advanced stage. More sophisticated tests such as magnetic resonance spectroscopy are more meaningful, but they are not widely applicable because of the relatively high costs in clinical practice. Therefore, scientists around the world are working on simple and accurate methods that are more suitable for diagnosing a fatty liver. The Fatty Liver Index (FLI) consisting of the parameters age, waist circumference, and triglyceride levels (TG) measured in the blood in a fasting state, and gamma-glutamyl transpeptidase (GGT), has been shown to be quite effective.

The researchers from Tübingen led by Professor Stefan, in collaboration with Professor Schulze of the German Institute of Human Nutrition (DIfE), a partner of the DZD, and their colleagues from Hamburg and Italy, have developed an improved FLI. For this purpose, in addition to the parameters of the FLI, they used the TG and glucose levels from an oral glucose tolerance test measured at the time point of two hours as well as the most important gene variant for fatty liver (rs738409 C>G in PNPLA3). On the basis of data from the TULIP (Tübingen Lifestyle Intervention Program) study, they showed that with this new and extended fatty liver index, diagnosis was improved over the standard FLI, and also more precisely determined the probability of reducing liver fat content during a lifestyle intervention. Professor Stefan now hopes that "this index will be increasingly used in clinical practice to diagnose fatty [liver disease](#) at an [early stage](#) in order to prevent the consequences of [fatty liver disease](#)."

More information: K. Kantartzis et al, An extended fatty liver index to predict non-alcoholic fatty liver disease, *Diabetes & Metabolism* (2017). DOI: [10.1016/j.diabet.2016.11.006](https://doi.org/10.1016/j.diabet.2016.11.006)

Provided by Deutsches Zentrum fuer
Diabetesforschung DZD

APA citation: A new index for the diagnosis of non-alcoholic fatty liver disease (2017, January 24)
retrieved 4 September 2022 from <https://medicalxpress.com/news/2017-01-index-diagnosis-non-alcoholic-fatty-liver.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.