

# Decreased physical activity common after cancer diagnosis

25 January 2017



improve many outcomes in patients with cancer," lead researcher Jun Mao, M.D., chief of integrative medicine service at the Memorial Sloan Kettering Cancer Center in New York City, told *HealthDay*. Doctors need to ask their patients about [physical activity](#) and encourage them to be active, he said.

**More information:** [Press Release](#)  
[More Information](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

(HealthDay)—Exercise can help cancer patients cope with their treatment, but as many as 75 percent reduce their physical activity after diagnosis, according to a study presented at the American Society of Clinical Oncology's Cancer Survivorship Symposium, held from Jan. 27 to 28 in San Diego.

The study included 662 [cancer patients](#) being treated at 12 Philadelphia clinics. Most of the patients (65 percent) were overweight or obese women. Their average age was 60.

Surveys revealed that 75 percent of patients said they had reduced their physical activity since being diagnosed with cancer. Sixteen percent said they had kept up the same level of activity as they had before diagnosis, and 4 percent said they had increased their exercise since diagnosis.

"We have about 25 million cancer survivors in the United States, and there is mounting evidence that suggests that physical activity and exercise

APA citation: Decreased physical activity common after cancer diagnosis (2017, January 25) retrieved 29 April 2021 from <https://medicalxpress.com/news/2017-01-decreased-physical-common-cancer-diagnosis.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*