

New guideline provides clinical recommendations for specific insomnia drugs

February 15 2017

A new clinical practice guideline is the first from the American Academy of Sleep Medicine to provide comprehensive, evidence-based analyses of individual agents commonly used in the treatment of chronic insomnia disorder.

The guideline, which is published in the Feb. 15 issue of the *Journal of Clinical Sleep Medicine*, provides recommendations to help clinicians choose a specific pharmacological agent for the treatment of chronic insomnia in adults, when such treatment is indicated. Developed by an expert task force and approved by the AASM board of directors, the guideline was based on a systematic literature review, meta-analyses, and assessment of the evidence using the GRADE methodology. A draft of the guideline was previously made available for public comment.

"The publication of this [clinical practice guideline](#) is an important step forward for the field of [sleep medicine](#)," said AASM President Ronald D. Chervin, MD, MS. "It further equips clinicians to provide high quality, patient-centered care for millions of people who suffer from chronic insomnia."

Although transient insomnia symptoms occur in 30 to 35 percent of the population, the full clinical syndrome of chronic insomnia disorder occurs in about 10 percent of people. The [CDC has estimated](#) that 4.1 percent of U.S. adults report having taken a prescription sleep aid in the past month, based on survey data collected from 2005 - 2010.

The [clinical practice](#) guideline includes 14 specific recommendations for individual agents, including prescription medications such as zolpidem and eszopiclone; over-the-counter medications such as diphenhydramine; and dietary supplements such as melatonin and valerian. Each recommendation suggests whether clinicians should or should not use the individual drug for sleep onset insomnia or sleep maintenance insomnia, versus no treatment. However, the guideline does not recommend one drug over another since few comparative efficacy studies have been conducted among these agents.

The authors noted that the data available to support these recommendations is often less than certain. As a result, the strength of each recommendation is classified as "weak," reflecting a lower degree of certainty in the appropriateness of the patient-care strategy. Therefore, clinicians must continue to exercise sound clinical judgment in their care for patients with chronic insomnia. Decisions should be based not only on these recommendations but also on a clinician's experience and the individual circumstances presented by the patient.

The guideline emphasizes that medications for chronic insomnia disorder should be considered mainly in patients who are unable to participate in cognitive behavioral therapy for insomnia (CBT-I), patients who still have symptoms after this therapy, or those who require a temporary adjunct to CBT-I. A [clinical guideline](#) previously published by the American Academy of Sleep Medicine indicates that CBT-I is an effective treatment that should be utilized as an initial intervention for [chronic insomnia](#), and this position was reiterated by the AASM in a 2014 [Choosing Wisely statement](#).

Anyone who is having trouble falling asleep or staying asleep should discuss the concern with a doctor. Help for an ongoing sleep problem is available from more than 2,500 sleep centers that are accredited by the American Academy of Sleep Medicine.

More information: Michael J. Sateia et al, Clinical Practice Guideline for the Pharmacologic Treatment of Chronic Insomnia in Adults: An American Academy of Sleep Medicine Clinical Practice Guideline, *Journal of Clinical Sleep Medicine* (2017). [DOI: 10.5664/jcsm.6470](https://doi.org/10.5664/jcsm.6470)

Provided by American Academy of Sleep Medicine

Citation: New guideline provides clinical recommendations for specific insomnia drugs (2017, February 15) retrieved 20 December 2022 from <https://medicalxpress.com/news/2017-02-guideline-clinical-specific-insomnia-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.