

Burnout is prevalent among pediatric residents

February 25 2017



(HealthDay)—Burnout is prevalent among pediatric residents and is

associated with suboptimal patient care attitudes and behaviors, according to research published online Feb. 23 in *Pediatrics*.

Tamara Elizabeth Baer, M.D., M.P.H., from Boston Children's Hospital, and colleagues surveyed 258 residents from 11 pediatric residency programs to assess the prevalence of [burnout](#). Using seven questions from a standardized qualitative survey the authors measured patient care attitudes and behaviors.

The researchers found that 39 percent of respondents endorsed burnout. The odds of reporting suboptimal patient care attitudes and behaviors were significantly elevated among residents with burnout; these included discharging patients to make the service more manageable, not fully answering questions or discussing treatment options, making treatment or [medication errors](#), ignoring the social or personal aspect of illness, and feeling guilty about patient treatment (adjusted odds ratios, 4.2, 3.5, 7.1, 9.6, and 6.0, respectively).

"Residency programs should develop interventions addressing burnout and its potential negative impact on [patient care](#)," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

[Editorial \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Burnout is prevalent among pediatric residents (2017, February 25) retrieved 9 January 2023 from <https://medicalxpress.com/news/2017-02-burnout-prevalent-pediatric-residents.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.