

# High cholesterol levels linked with rotator cuff surgery failure

14 March 2017

Patients with higher cholesterol levels face a significantly greater risk for failure of minimally invasive (arthroscopic) rotator cuff surgery, according to a new study to be presented on Friday, March 17, at the 2017 Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS). According to the study, commonly-used cholesterol lowering drugs, called statins, diminish this risk.

"The study found an association between the use of statin lipid lowering agents and mitigation of the risk for revision surgery, even in patients with elevated [cholesterol](#) levels," said Brian C. Werner, MD, an orthopaedic surgeon at the University of Virginia School of Medicine and lead author of the study.

Approximately 200,000 Americans undergo [rotator cuff](#) surgery, the repair of the primary tendons and muscles in the shoulder, each year.

In the study, researchers reviewed a private insurer patient database of nearly 31,000 patients undergoing arthroscopic rotator cuff repair from 2007 to 2014. Patients were divided into normal, moderate and high cholesterol groups, based on their total cholesterol levels (the measure of low-density lipoprotein [LDL], also known as "bad" cholesterol, and high-density lipoprotein [HDL], or "[good cholesterol](#)"), LDL levels alone, and triglycerides, another form of fat in the blood that increases the risk of heart disease. The patient groups were further analyzed based on whether or not they took statins before and after surgery.

Among the findings of the research:

- The rate of revision [rotator cuff surgery](#) was significantly increased in patients with moderate or high total cholesterol and LDL levels compared to patients with normal levels.
- Among the patients with moderate or high

cholesterol, those who were not taking statins before or after surgery had significantly higher rates of revision surgery.

- Patients taking statins prior to or after surgery did not have higher rates of revision surgery, even if their [cholesterol levels](#) were high.
- Patient triglyceride levels had no impact on rotator cuff [surgery](#) outcomes.

"Additional study of the association between statins and rotator cuff surgical outcomes is encouraged to determine if improved cholesterol control can improve clinical outcomes following arthroscopic rotator cuff repair," said Dr. Werner.

**More information:** Study abstract: [submissions.miramsmart.com/Veri ...  
on/temp/radC7F94.pdf](https://submissions.miramsmart.com/Veri...on/temp/radC7F94.pdf)

Provided by American Academy of Orthopaedic Surgeons

APA citation: High cholesterol levels linked with rotator cuff surgery failure (2017, March 14) retrieved 4 May 2021 from <https://medicalxpress.com/news/2017-03-high-cholesterol-linked-rotator-cuff.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*