

# Eating more fruits and vegetables may lower risk of blockages in leg arteries

18 May 2017



Credit: Anna Langova/Public Domain

Eating three or more servings of fruit and vegetables per day may lower your risk of developing peripheral artery disease (PAD), according to new research in *Arteriosclerosis, Thrombosis and Vascular Biology*, an American Heart Association journal.

PAD narrows the arteries of the legs, limiting blood flow to the muscles and making it difficult or painful to walk or stand.

Previous studies linked lower consumption of fruits and vegetables with the increased occurrence of [coronary heart disease](#) and stroke. However, there has been little research into the association of eating fruits and vegetables and PAD.

After studying data from 3.7 million people, researchers found:

- People who reported eating three or more daily servings of fruits and vegetables had 18 percent lower odds of PAD than those

reporting eating less.

- When stratified by smoking status, the association of lower PAD and increased fruits and vegetables was present only among participants who were current or former smokers.
- Overall, 6.3 percent of participants had PAD and 29.2 percent reported eating three or more servings of fruits and vegetables daily.

"Our current study provides important information to the public that something as simple as adding more fruits and vegetables to your diet could have a major impact on the prevalence of life-altering [peripheral artery disease](#)," said Jeffrey Berger, M.D., study coauthor and associate professor of medicine and surgery at New York University School of Medicine in New York City.

Participants, who were average age 64 (64 percent women, nearly 90 percent white), completed medical and lifestyle questionnaires and [ankle brachial index](#) tests at more than 20,000 sites across America. An ankle brachial index test is a comparison of blood pressure differences between readings at the ankle and the forearm.

Researchers also said their study confirmed that Americans' overall fruit and [vegetable intake](#) remains dismally low.

The association of fruit and vegetable intake and lower PAD risk persisted after accounting for age, gender, race, smoking status and multiple other cardiovascular risk factors. Researchers noted older white women were most likely to eat three or more servings of fruits and vegetables daily, while younger black men were the least likely to report daily intake of three or more servings of fruits and vegetables. Low fruit and vegetable intake was particularly associated with PAD among current and former smokers.

"Our study gives further evidence for the

importance of incorporating more fruits and vegetables in the diet," said Sean Heffron, M.D., M.S., M.Sc., study coauthor and instructor in medicine at New York University School of Medicine. "One-on-one dietary assessments and counseling for PAD patients, as well as greater public health awareness of the importance of [fruit](#) and [vegetable](#) consumption, are both needed."

**More information:** *Arteriosclerosis, Thrombosis and Vascular Biology* (2017). [DOI: 10.1161/ATVBAHA.116.308474](#)

Provided by American Heart Association

APA citation: Eating more fruits and vegetables may lower risk of blockages in leg arteries (2017, May 18) retrieved 7 July 2022 from <https://medicalxpress.com/news/2017-05-fruits-vegetables-blockages-leg-arteries.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*