

Investigators examine the relationship between pain and opioid abuse

12 June 2017

The drug overdose epidemic is largely driven by opioids, which continue to be prescribed for chronic pain despite recommendations to use non-opioids for most cases. A new review published in the *British Journal of Pharmacology* examines the interaction between pain and the abuse of opioids, and investigates the circuits in the brain that may be behind this link. The review is part of a special theme issue on Emergent Areas of Opioid Pharmacology.

"We have shown that the brain's natural [opioid](#) system is drastically changed by the presence of pain, and these changes may very well contribute to the difficulty of treating chronic pain with opioids," said first author Adrienne Wilson-Poe, PhD of the Washington University in Saint Louis School of Medicine. "We have just glimpsed the tip of the iceberg when it comes to pain's effect on the brain, however, and we need a lot more research and grant funding to get to the bottom of the extremely complex interaction between drug abuse and pain."

She and senior author Jose Moron-Concepcion, PhD, Associate Professor in the Department of Anesthesiology at Washington University, note that without a fundamental understanding of pain-induced changes in the brain and how these adaptations interact with subsequent drug exposure, investigators are merely fishing for solutions to the opioid crisis. "Our work is attacking this problem head-on by diligently characterizing the mechanisms involved in pain, addiction, and the interaction between them," said Dr. Wilson-Poe. "We envision a future where chronic pain is considered a disease in its own right, not merely a symptom of some other biological process."

The review stresses that opioids are the most powerful analgesics known to man, and their continued use in the treatment of severe [pain](#) is inevitable; however, [opioid therapy](#) of the future must look very different from how it does today.

Efforts to address this issue include a 2016 guideline by the Centers for Disease Control and Prevention that recommends using non-opioids for most cases of [chronic pain](#), using the lowest effective dose when prescribing opioids, and ensuring that patients who are treated with opioids are closely monitored.

The review is part of a larger themed issue, 'Emergent Areas of Opioid Pharmacology,' that will publish at a later time.

The National Institute on Drug Abuse notes that the emergence of illicitly manufactured synthetic opioids including fentanyl, carfentanil, and their analogues represents an escalation of the ongoing opioid overdose epidemic. Also, prescription opioid misuse is a significant risk factor for heroin use, and 80% of heroin users first misuse [prescription opioids](#).

More information: Adrienne R Wilson-Poe et al, The dynamic interaction between pain and opioid misuse, *British Journal of Pharmacology* (2017). DOI: [10.1111/bph.13873](https://doi.org/10.1111/bph.13873)

Provided by Wiley

APA citation: Investigators examine the relationship between pain and opioid abuse (2017, June 12)
retrieved 27 April 2021 from
<https://medicalxpress.com/news/2017-06-relationship-pain-opioid-abuse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.