

Gap in health care for Alzheimer's disease patients who live alone

June 15 2017




Diagram of the brain of a person with Alzheimer's Disease. Credit: Wikipedia/public domain.

Swedish researchers report in an article published in the *Journal of Alzheimer's Disease* that 46 percent of patients who are diagnosed with Alzheimer's disease in Sweden live alone in their homes, in particular older women.

The patients who live alone do not receive the same extent of diagnostic investigations and anti-dementia treatment as those who are co-habiting. On the other hand, they were treated more frequently with antidepressants, antipsychotics and sedative drugs.

According to recent statistics, the number of older people who live alone in their homes, especially women, is increasing in high income countries. When an older person is affected by dementia, such as [Alzheimer's disease](#), they may not have a close relative living with them, which may complicate the course of the disease. Dementia affects their memory and later can lead to their dependency on caregivers.

Investigators at Karolinska Institutet and Karolinska University Hospital in Sweden studied 26 000 patients with Alzheimer's disease registered in the Swedish Dementia Registry (SveDem), a nationwide quality register on patients affected by dementia. Owing to the size of the study sample and detailed information received from The Swedish Prescribed Drug Register and The Swedish Patient Register, the team was able to estimate the effects of solitary living on several aspects of [health care](#), independently of patients' characteristics, such as age, gender, comorbidities or severity of dementia.

"The high proportion of solitary living Alzheimer's disease patients is worrying as this is certainly not a disease patients can deal with alone," says Pavla Čermáková, the first author of the study, on behalf of the team from the Department of Neurobiology, Care Sciences and Society at Karolinska Institutet. "We found many inequalities in the health care that is provided. Solitary living patients received less frequently investigations with brain imaging methods, such as computed tomography (CT) and [magnetic resonance imaging](#) (MRI), and lumbar puncture. Moreover, they were treated less commonly with cholinesterase inhibitors and memantine, which are nowadays the only drugs that can improve their symptoms," says dr. Čermáková.

A likely explanation is that the presence of a close caregiver who lives with the patient influences how physicians decide about the diagnostic strategy and therapy. On the contrary, patients who live alone got more antidepressants, antipsychotics and hypnotics-sedatives. It is not clear

whether the doctors are more prone to prescribe these drugs to this patient group or if solitary living can cause or worsen depression, sleeping problems and psychotic symptoms.

"It is possible that improving living conditions of patients with Alzheimer's disease could lower the need for these drugs," suggests Maja Nelson, who is the second author of the paper. This study provides a good basis for a possible intervention. "I work clinically at the Karolinska University Hospital where I see patients with Alzheimer's disease," says Dorota Religa, who is the senior author of the study. "We need to make more efforts to overcome challenges that solitary living poses on them," adds Dr. Religa.

Provided by IOS Press

Citation: Gap in health care for Alzheimer's disease patients who live alone (2017, June 15) retrieved 23 December 2023 from <https://medicalxpress.com/news/2017-06-gap-health-alzheimers-disease-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.