

Shingles increases risk of heart attack, stroke

3 July 2017

Contracting shingles, a reactivation of the chickenpox virus, increases a person's risk of stroke and heart attack, according to a research letter published today in the *Journal of the American College of Cardiology*.

According to the Centers for Disease Control and Prevention, almost 1 out of every 3 people in the United States will develop shingles in their lifetime. Anyone who has suffered from chickenpox may develop shingles; however, the risk of shingles increases as a person gets older.

Researchers in South Korea used the National Health Insurance Service's "medical check-up" database to identify <u>patients</u> with newly diagnosed herpes zoster—or shingles, stroke and <u>heart attack</u> using the relevant International Classification of Disease-10 diagnostics codes.

A total of 519,880 patients were followed from 2003-2013, during this period there were 23,233 cases of shingles. The final cohort of 23,213 was matched with the same number of shingles-free patients to serve as control subjects.

Patients with shingles were more likely to be female and common risk factors for stroke and <u>heart</u> attack, such as old age, <u>high blood pressure</u>, diabetes and high cholesterol, were also more commonly seen in these patients. However, this group was also less likely to smoke, have a lower alcohol intake, more exercise and be part of a higher socioeconomic class.

Shingles was found to raise the risk of a composite of cardiovascular events including heart attack and stroke by 41 percent, the risk of stroke by 35 percent and the risk of heart attack by 59 percent. The risk for stroke was highest in those under 40 years old, a relatively younger population with fewer risks for atherosclerosis. The risks of both stroke and heart attack were highest the first year after the onset of shingles and decreased with

time. However, these risks were evenly distributed in the shingles-free group.

"While these findings require further study into the mechanism that causes <u>shingles</u> patients to have an increased risk of heart attack and <u>stroke</u>, it is important that physicians treating these patients make them aware of their increased risk," said Sung-Han Kim, MD, PhD, a physician in the department of infectious diseases at Asan Medical Center in Seoul and one of the study authors.

Provided by American College of Cardiology



APA citation: Shingles increases risk of heart attack, stroke (2017, July 3) retrieved 4 December 2022 from <u>https://medicalxpress.com/news/2017-07-shingles-heart.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.