


# Aphasia recovery via speech therapy related to structural plasticity of the ventral stream

July 10 2017



Pre- and post-treatment MK values along the core ILF from a representative participant are shown anatomically. The ILF bottleneck, which is marked with a black arrow, demonstrates an increase in MK toward normal values from before to after therapy. This participant demonstrated a 55 percent improvement in semantic errors. Credit: Dr. Leonardo Bonilha and Emilie McKinnon. All rights reserved.

Language disturbances (aphasia) are common after stroke and can manifest as difficulty identifying the correct word to use (semantic problems) and/or difficulty pronouncing words (phonemic problems). Speech therapy has long been a standard of post-stroke care and can improve aphasia. However, the neurological basis of therapy-mediated recovery is poorly understood, and researchers do not know why some patients improve with therapy while others show little response.

The ability of the residual language network to rebuild itself (a characteristic called structural plasticity) is directly related to how much benefit a patient receives from speech therapy, report investigators at the Medical University of South Carolina (MUSC) in an article published online June 19, 2017 by *Annals of Neurology*. Their study also supports the dual-stream language model, which holds that ventral brain networks are associated with semantic skills and dorsal networks with phonemic skills.

"Producing speech is a two-step process - first selecting the correct word using semantic associations and, then, pronouncing it phonetically," explains lead author Emilie T. McKinnon, an M.D., PhD candidate in MUSC's Department of Neurology. "The current theory is that different parts of the brain house these two processes. If that's the case, these areas have to communicate with each other to produce language. So, the question is, when one of those regions is damaged, how is that connection restored? We looked at the microstructure of one of those connections to try to see what happens when language processing improves."

The team tested eight aphasia patients, all of whom had had a single stroke at least one year prior that affected the ILF region. Participants were tested for confrontational naming ability one week before and one week after receiving a three-week course of intensive speech therapy. In addition, participants underwent four [magnetic resonance imaging](#)

(MRI) sessions - two before speech therapy and two after.

A novel aspect of the study was how the MRIs were conducted and assessed. The investigators leveraged recent advancements in diffusion-weighted imaging and image analysis, providing greater sensitivity to microstructural white matter changes and revealing previously hidden differences.

"Diffusion-weighted imaging has been around a long time and it's useful for mapping and studying brain networks, but the commonly used analysis techniques can be improved upon," explains McKinnon. "We used diffusion kurtosis imaging and kurtosis-based tractography to calculate how water diffuses in the brain and identify areas of high resistance. Where water diffuses unimpeded, kurtosis is near zero. The more barriers it meets, the higher the kurtosis. So, it's a measure of how complex the environment is. We used this strategy because we could get so much more information and it only required adjusting the MRI settings and adding about five minutes of scanner time."

The team focused in each patient on the ILF area with the lowest mean kurtosis (MK), indicating that this segment, the 'bottleneck', had the most damage. They found a significant correlation between pre- to post-therapy MK increases at the bottleneck (i.e., kurtosis values changing toward normal ranges in the most damaged area) and therapy-related semantic [language](#) improvements ( $r=-0.90$ ,  $p$

Citation: Aphasia recovery via speech therapy related to structural plasticity of the ventral stream (2017, July 10) retrieved 15 July 2023 from <https://medicalxpress.com/news/2017-07-aphasia-recovery-speech-therapy-plasticity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.