

AAIC: Turnover kinetics vary for different amyloid beta isoforms

July 20 2017



(HealthDay)—Amyloid beta ($A\beta$)₃₈ has faster turnover kinetics than

A β 40 and A β 42, according to a study published online July 19 in *Alzheimer's & Dementia* to coincide with presentation at the annual Alzheimer's Association International Conference, held from July 16 to 20 in London.

Vitaliy Ovod, from the Washington University School of Medicine in St. Louis, and colleagues examined the turnover kinetics and concentrations of A β 38, A β 40, and A β 42 in [human plasma](#) using the Stable Isotope Labeling Kinetics protocol.

The researchers found that the half-life of A β isoforms was approximately three hours in [plasma](#). Compared with A β 40 and A β 42, A β 38 demonstrated faster turnover kinetics. Amyloid positive participants had faster fractional turnover of A β 42/A β 40 and lower A β 42 and A β 42/A β 40 concentrations.

"Faster turnover kinetics and lower A β 42 concentrations in amyloid positive participants have been seen in prior cerebrospinal fluid A β studies, suggesting a blood-brain transportation mechanism of A β ," the authors write. "The stability and sensitivity of plasma A β measurements [suggest] this may be a useful screening test for central nervous system amyloidosis."

More information: [Abstract](#)

[Full Text \(subscription or payment may be required\)](#)

[More Information](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: AAIC: Turnover kinetics vary for different amyloid beta isoforms (2017, July 20) retrieved 19 November 2023 from

<https://medicalxpress.com/news/2017-07-aaic-turnover-kinetics-vary-amyloid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.