

Moderate, vigorous activity not tied to more elderly falls

3 August 2017



the previous year (interaction P falls in women with lower MVPA levels (13.0 percent; P

"These findings support encouraging [women](#) to engage in the amounts and types of MVPA that they prefer," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

(HealthDay)—Falls are not more common or injurious in older women who engage in higher levels of moderate-to-vigorous physical activity (MVPA), according to a study published online July 29 in the *Journal of the American Geriatrics Society*.

David M. Buchner, M.D., from the University of Illinois at Urbana-Champaign, and colleagues assessed whether MVPA (measured using an accelerometer) is associated with incident falls among 5,545 ambulatory women aged 63 to 99 years participating in the Women's Health Initiative.

The researchers observed a greater fall risk in women in the lowest quartile of MVPA versus those in the highest (incident rate ratio, 1.18), after adjusting for age, race and ethnicity, and fall risk factors. Fall rates were significantly associated with MVPA in women with low Short Physical Performance Battery (SPPB) scores (?8) or a history of frequent falls, but not in women with high SPPB scores (9 to 12) or no more than one fall in

APA citation: Moderate, vigorous activity not tied to more elderly falls (2017, August 3) retrieved 17 August 2022 from <https://medicalxpress.com/news/2017-08-moderate-vigorous-tied-elderly-falls.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.