

Subcutaneous exendin treats post-bariatric hypoglycemia

August 11 2017



(HealthDay)—Subcutaneous exendin (SC Ex-9) appears to be safe and

effective in treating post-bariatric hypoglycemia (PBH), according to a study published online Aug. 4 in *Diabetes, Obesity and Metabolism*.

Colleen M. Craig, M.D., from the Stanford University School of Medicine in California, and colleagues conducted the first in-human subcutaneous administration of Ex-9 in nine female patients with PBH following Roux-en-Y gastric bypass. First, a single participant underwent equimolar low-dose IV versus subcutaneous Ex-9 administration. Then, eight participants were administered single ascending doses of SC Ex-9 during [oral glucose tolerance](#) testing.

The researchers observed an exposure-response relationship, but all doses effectively prevented hyperinsulinemic hypoglycemia and improved associated symptoms. The postprandial glucose nadir was increased by 66 percent, peak insulin was reduced by 57 percent, and neuroglycopenic symptoms were reduced by 80 percent, on average. No treatment-related adverse events were reported.

"SC Ex-9 appears to represent a safe, effective, and targeted therapeutic approach for treatment of PBH," the authors write. "Further investigation involving multiple doses with chronic dosing is warranted."

Several authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Subcutaneous exendin treats post-bariatric hypoglycemia (2017, August 11) retrieved 23 April 2023 from <https://medicalxpress.com/news/2017-08-subcutaneous-exendin-post-bariatric->

[hypoglycemia.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.