

Potential treatment for brain cancer as drug shrinks tumours

14 August 2017



L-R: Dr Amanda Millar, Dr Laura Genovesi and Professor Brandon Wainwright. Credit: University of Queensland

An international team of researchers has found a drug previously approved to treat breast cancer could also be used to shrink medulloblastoma, a common form of childhood brain tumour.

The discovery, made by The University of Queensland's Institute for Molecular Bioscience and the Fred Hutchinson Cancer Research Center in Seattle, has led to a clinical trial using the drug palbociclib to treat children with <u>medulloblastoma</u>, the most common malignant brain <u>tumour</u> found in children.

UQ's Professor Brandon Wainwright said <u>brain</u> <u>tumours</u> were the most common cause of cancer death in infants, children and adolescents – and even survivors can end up with significant longterm side effects from existing treatments.

"Clearly, we need new therapies that increase survival of young patients and reduce the side effects they suffer, such as delays in brain development, growth problems and increased risk of other cancers," Professor Wainwright said. Fellow UQ researcher Dr Laura Genovesi examined the genetic code of medulloblastoma to predict whether these tumours may respond to alreadyapproved drugs.

"This analysis led us to believe that palbociclib, an oral <u>drug</u> approved in 2015 for the treatment of <u>breast cancer</u>, would be effective against medulloblastoma," Dr Genovesi said.

"We expected that palbociclib would arrest the growth of medulloblastoma, but we were stunned to find that it went a step further and actually shrank the tumours to a size where survival is possible."

"The finding is remarkable since the tumours were very advanced and were treated for only a short period of time and we did not use any other therapy such as chemotherapy in combination.

"Some tumours recurred once treatment with palbociclib stopped, probably due to resistant cells within the tumour.

"It means that palbociclib, or drugs like it, could be used against medulloblastoma in combination with other drugs to treat resistant cells.

"If the clinical trial is successful, it would represent a major step forward to taking this research from the genome to the clinic."

The study is published in the journal *Clinical Cancer Research*.

More information: Michelle L Cook Sangar et al. Inhibition of CDK4/6 by Palbociclib Significantly Extends Survival in Medulloblastoma Patient-Derived Xenograft Mouse Models, *Clinical Cancer Research* (2017). <u>DOI:</u> <u>10.1158/1078-0432.CCR-16-2943</u>



Provided by University of Queensland

APA citation: Potential treatment for brain cancer as drug shrinks tumours (2017, August 14) retrieved 10 June 2022 from <u>https://medicalxpress.com/news/2017-08-potential-treatment-brain-cancer-drug.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.