




# Key synapse formation regulator identified

22 August 2017



is also known that the MDGA1 protein binds neuroigin-2 at the inhibitory [synapses](#) and interferes with the interaction of neuroigin and neurexin and stops inhibitory synapse development, but its [mechanism](#) is also unclear.



Molecular Structure of Synapse Regulatory Protein binding Neuroigin-2 and MDGA1. Through protein crystallography, researchers were able to derive a key region that plays a pivotal role in the binding and function of the two proteins. Credit: DGIST

Neuroigin-2 /MDGA1 and Neuroigin-1 / Neurexin1β binding sites partially overlap and researchers proposed a binding competition model between MDGA1 and Neurexin1β in this part. Credit: DGIST


Professor Ko Jae-won at Korea Advanced Institute of Science and Technology (KAIST) has conducted a study of the three-dimensional structure of proteins that regulate neuronal cell connections for the first time, and has identified the control mechanisms of synapse formation.

The research teams used protein crystallography to crystallize neuroigin-2 and MDGA1 protein complexes that are involved in inhibitory synapse development and observed the [three-dimensional structure](#). The researchers have identified for the first time that the MDGA1 protein at the post-synapse interferes with the binding between neuroigin-2 and neurexin and prevents inhibitory synapse formation.

The brain is made up of innumerable [neurons](#). Neurotransmitters are secreted and absorbed at the synapse, the junction of the two neurons; the signals are transmitted and the brain functions properly. As the brain develops, efficient neurotransmission relies on a balance of [excitatory synapses](#) and inhibitory synapses created between the neurons. Excitatory synapses release neurotransmitters that excite neurons and synaptic inhibitors release neurotransmitters to inhibit nerve cell excitability. When the balance of the two synapses is broken, it is known that mental disorders such as autism, bipolar disorder and obsessive-compulsive disorder occur.

Based on the three-dimensional molecular structure, the researchers created mutations of core amino acids that play a pivotal role in the binding of neuroigin-2 and MDGA1. Through the neuron culture experiment, they verified that the protein interaction site is a critical functional region for the negative regulation of the synapse development process. In addition, the study has proved that MDGA1 can effectively control inhibitory synapse formation as the binding ability of MDGA1 is superior to neurexin while MDGA1 and [neurexin](#) can both competitively bind with neuroigin-2.

In 2013, Professor Thomas C. Sudhof of Stanford University discovered the neuroigin and neurexin proteins, important synaptic adhesion proteins that are important factors in synapse development and function maintenance. However, specific mechanisms have yet to be identified. Moreover, it



Neurexins (orange) at the presynapse and Neuroligin at the postsynapse mediate synaptic development by binding two neurons through interactions. In inhibitory synapses, the regulatory protein MDGA1 produces a Neuroligin-2 (NL2) / MDGA1 complex based on strong binding power to Neuroligin-2 / Neurexin and effectively inhibits the development of inhibitory synapse by selectively interfering with binding of Neuroligin-2/Neurexin. Credit: DGIST

DGIST's Professor Jaewon Ko of the Department of Brain and Cognitive Sciences said, "We have found the mechanism of molecular regulation of MDGA1 [protein](#), which is necessary for the balanced operation of excitatory and [inhibitory synapses](#). We will continue working to identify the mechanisms of brain diseases caused by dysfunction of synaptic proteins and conduct researches to develop therapeutic drugs."

This study has been published in the online edition of *Neuron*, an international journal of neurobiology and the sister journal of *Cell*, on June 21.

**More information:** Jung A Kim et al, Structural Insights into Modulation of Neurexin-Neuroligin Trans -synaptic Adhesion by MDGA1/Neuroligin-2 Complex, *Neuron* (2017). [DOI: 10.1016/j.neuron.2017.05.034](https://doi.org/10.1016/j.neuron.2017.05.034)

Provided by DGIST

APA citation: Key synapse formation regulator identified (2017, August 22) retrieved 3 June 2022 from <https://medicalxpress.com/news/2017-08-key-synapse-formation.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*