


Cancer drug may benefit patients with inherited form of kidney disease

24 August 2017



Credit: CC0 Public Domain

A cancer drug called bosutinib may inhibit the growth of cysts in patients with autosomal dominant polycystic kidney disease (ADPKD), according to a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*. The findings point to a potential new treatment strategy for affected patients, but the long-term benefits remain to be determined.

ADPKD is an inherited disorder that affects up to 1 in 1000 people and is characterized by cysts in the [kidney](#) and other organs. As patients' kidney volume increases due to cyst growth, they gradually lose their [kidney function](#) and often develop [kidney failure](#). Current treatments are primarily supportive, such as focusing on hypertension and other secondary complications.

The inherited mutations that cause ADPKD affect a protein involved in various signaling pathways that often involve enzymes called [tyrosine kinases](#). Therefore, a team led by Vladimir Tesar, MD, PhD (Charles University and General University Hospital, in the Czech Republic) tested the potential of an investigational drug called bosutinib

that inhibits a particular tyrosine kinase called Src/Bcr-Abl.

The phase 2 study included patients with ADPKD who were randomized 1:1:1 to bosutinib 200 mg/day, bosutinib 400 mg/day, or placebo. Of 172 patients enrolled, 169 received at least one treatment. The higher dose of bosutinib was not well tolerated.

The annual rate of kidney enlargement was reduced by 66% for patients receiving bosutinib 200 mg/day vs. those receiving placebo (1.63% vs. 4.74%, respectively) and by 82% for all patients receiving bosutinib vs. those receiving placebo (0.84% vs. 4.74%, respectively). The study was not powered to demonstrate a treatment effect on kidney function, but there was no evidence of a benefit associated with bosutinib compared with placebo over the 2-year treatment period.

"The reduction in growth of cysts through [treatment](#) with bosutinib was confirmed, although gastrointestinal side effects (primarily diarrhea), which were partly dose-dependent, may represent a substantial drawback for the further development of the drug for [patients](#) with ADPKD," said Prof. Tesar.

More information: "Bosutinib Versus Placebo for Autosomal Dominant Polycystic Kidney Disease," *Journal of the American Society of Nephrology* (2017). [DOI: 10.2215/CJN.01530217](https://doi.org/10.2215/CJN.01530217)

Provided by American Society of Nephrology

APA citation: Cancer drug may benefit patients with inherited form of kidney disease (2017, August 24) retrieved 19 July 2022 from <https://medicalxpress.com/news/2017-08-cancer-drug-benefit-patients-inherited.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.