

Two-stage approach to risk-reducing mastectomy improves results for women with large breasts

29 August 2017

For women undergoing risk-reducing mastectomy to prevent breast cancer, reconstruction can be challenging in those with larger breasts. A two-stage approach—with initial breast reduction and "pre-shaping" followed by mastectomy and reconstruction—appears to be a safer procedure with better cosmetic results, reports the September issue of *Plastic and Reconstructive Surgery*, the official medical journal of the American Society of Plastic Surgeons (ASPS).

Targeted [breast](#) reduction/pre-shaping "primes" the breast for mastectomy and implant reconstruction performed a few months later, according the study by Gudjon L. Gunnarsson, MD, and colleagues of Telemark Hospital, Skien, Norway.

Breast Pre-Shaping Lowers Risks, Improves Final Results

More women are undergoing mastectomy and [breast reconstruction](#) to reduce their risk of familial [breast cancer](#), based on detection of specific risk genes. Typically this is done in a single-stage procedure, with "nipple-sparing" mastectomy of both breasts followed by immediate breast reconstruction using implants.

However, it can be challenging to achieve good results with immediate reconstruction in women with large breasts (macromastia) with sagging or drooping (ptosis). These patients "more often end up having a difficult corrective procedure and unacceptably high rate of failed reconstruction," Dr. Gunnarsson and coauthors write.

To improve their results, the researchers developed a targeted, two-stage approach to this group of patients. In the first stage, the women underwent a breast reduction/breast lift (mastopexy) procedure to create "a breast better

suited for implant reconstruction." In addition to reducing and reshaping the breast, an important part of this procedure was to lift the nipple-areola complex (nipple and surrounding tissues) to a more central position.

After a few months for healing, the women proceeded to the standard risk-reducing mastectomy and implant-based reconstruction. The online version of the article on the *Plastic and Reconstructive Surgery* website includes a video illustrating key aspects of the technique for plastic surgeons.

Between 2013 and 2015, Dr. Gunnarsson and colleagues used this two-stage approach in 22 women, median age 46 years. Mastectomy and subsequent reconstruction were successfully completed in all 44 breasts. The reconstructions were performed three to nine months (median four months) after initial breast reduction/reshaping.

Two patients had complications requiring repeat surgery; another five patients had minor complications that resolved with nonsurgical treatment. Importantly, there were no problems with the blood supply and survival of the nipple-areola complex.

In all cases, the cosmetic outcomes were rated excellent. The authors believe their technique is an important contribution to improving the results of risk-reducing mastectomy and [breast reconstruction](#) for women with large breasts. They note that the initial breast reduction/preshaping step "preconditions" the blood supply to the later reconstructed breast.

Dr. Gunnarsson and colleagues add that the two-stage procedure has become their "standard of care" for [women](#) with large breasts undergoing risk-

reducing [mastectomy](#). The researchers write, "The superior cosmetic outcome and low complication rate have proved to be better than we had anticipated."

More information: Gudjon L. Gunnarsson et al. Prophylactic Nipple-Sparing Mastectomy and Direct-to-Implant Reconstruction of the Large and Ptotic Breast, *Plastic and Reconstructive Surgery* (2017). [DOI: 10.1097/PRS.0000000000003621](https://doi.org/10.1097/PRS.0000000000003621)

Provided by Wolters Kluwer Health

APA citation: Two-stage approach to risk-reducing mastectomy improves results for women with large breasts (2017, August 29) retrieved 25 October 2022 from <https://medicalxpress.com/news/2017-08-two-stage-approach-risk-reducing-mastectomy-results.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.