

Recommendations vary for vision screening in young children

5 September 2017

The U.S. Preventive Services Task Force (USPSTF) recommends vision screening at least once in all children 3 to 5 years of age to detect amblyopia (also known as "lazy eye") or its risk factors (a B recommendation); and concludes that the current evidence is insufficient to assess the balance of benefits and harms of vision screening in children younger than 3 years (an I statement). The report appears in the September 5 issue of *JAMA*.

A B recommendation indicates that there is high certainty that the net benefit is moderate, or there is moderate certainty that the net benefit is moderate to substantial. An I statement indicates that evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.

One of the most important causes of [vision](#) abnormalities in [children](#) is [amblyopia](#). Amblyopia is an alteration in the visual neural pathway in a child's developing brain that can lead to permanent vision loss in the affected eye. Among children younger than 6 years, 1 percent to 6 percent have amblyopia or its [risk factors](#). Early identification of vision abnormalities could prevent the development of amblyopia. To update its 2011 recommendation, the USPSTF reviewed the evidence on the accuracy of vision screening tests and the benefits and harms of vision screening and treatment.

The USPSTF is an independent, volunteer panel of experts that makes recommendations about the effectiveness of specific preventive care services such as screenings, counseling services, and preventive medications.

Detection

The USPSTF found adequate evidence that vision screening tools are accurate in detecting vision abnormalities, including refractive errors,

strabismus (a misalignment of the eyes) and amblyopia, and found inadequate evidence to compare screening accuracy across age groups (younger than 3 vs 3 years and older). Many studies of clinical accuracy did not enroll children younger than 3 years.

Benefits of Early Detection and Treatment

The USPSTF found adequate evidence that treatment of amblyopia or its risk factors in children ages 3 to 5 years leads to improved visual acuity and found inadequate evidence that treatment of amblyopia or its risk factors in children younger than 3 years leads to improved vision outcomes (i.e., visual acuity) or other benefits.

Harms of Early Detection and Treatment

The USPSTF found adequate evidence to assess harms of vision screening tests in children ages 3 to 5 years, including higher false-positive rates in low-prevalence populations. False-positive [screening](#) results may lead to overdiagnosis or unnecessary treatment. The USPSTF found adequate evidence to bound the potential harms of [vision screening](#) and treatment in children ages 3 to 5 years as small, based on the nature of the interventions, and found inadequate evidence on the harms of treatment in children younger than 3 years.

More information: *JAMA* (2017). jamanetwork.com/journals/jama/.../1001/jama.2017.11260

Provided by The JAMA Network Journals

APA citation: Recommendations vary for vision screening in young children (2017, September 5) retrieved 1 May 2021 from <https://medicalxpress.com/news/2017-09-vary-vision-screening-young-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.