


Certain older adults don't get to the hospital soon enough when experiencing a heart attack

18 October 2017



More information: *Journal of the American Geriatrics Society* (2017). [DOI: 10.1111/jgs.15102](https://doi.org/10.1111/jgs.15102)

Provided by Wiley

Credit: CC0 Public Domain

For individuals experiencing a heart attack, delays in getting to the hospital can have life-threatening consequences. A new study in the *Journal of the American Geriatrics Society* found that certain factors—non-white race, atypical symptoms, and heart failure—are linked with such delays in older individuals.

The study included 2500 patients aged 75 or older hospitalized for heart attack. Pre-hospital delay (six or more hours before getting to the hospital) was much more common (42%) than in studies of younger [heart attack](#) populations, in whom the reported prevalence ranges from 20% to 25%.

"Delays in presentation can have huge consequences for older adults with heart attacks. Based on the results of our study, we need to develop better clinical and [public health strategies](#) to ensure timely presentation, especially among non-white communities, those with [atypical symptoms](#), and those with heart failure," said lead author Dr. Gregory Ouellet, of Yale University.

APA citation: Certain older adults don't get to the hospital soon enough when experiencing a heart attack (2017, October 18) retrieved 11 October 2022 from <https://medicalxpress.com/news/2017-10-older-adults-dont-hospital-experiencing.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.