

Certain factors are linked to sleep disturbance in patients with diabetes

8 November 2017

In a *Journal of Advanced Nursing* study of 90 adults with type 2 diabetes, female gender, elevated blood sugar levels, neuropathic pain, and fatigue were related to sleep disturbance, even after controlling for factors such as age, diabetes duration, depressive symptoms, and distress.

The findings suggest that clinicians should pay extra attention to females when conducting sleep-related assessments. A comprehensive sleep evaluation should also include diabetes-related symptoms such as [neuropathic pain](#) and fatigue.

"Sleep disturbance in patients with diabetes is common and may negatively affect blood glucose," said senior author Dr. Cynthia Fritschi, of the University of Illinois at Chicago College of Nursing. "A thorough sleep assessment, especially in female adults, must include a symptom assessment. To do otherwise limits our ability to treat sleep disturbance effectively in this population."

More information: *Journal of Advanced Nursing* (2017). [DOI: 10.1111/jan.13482](https://doi.org/10.1111/jan.13482)

Provided by Wiley

APA citation: Certain factors are linked to sleep disturbance in patients with diabetes (2017, November 8) retrieved 30 May 2022 from <https://medicalxpress.com/news/2017-11-factors-linked-disturbance-patients-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.