

Small but distinct differences among species mark evolution of human brain

23 November 2017



Credit: Human Brain Project

The most dramatic divergence between humans and other primates can be found in the brain, the primary organ that gives our species its identity.

However, all regions of the human <u>brain</u> have molecular signatures very similar to those of our primate relatives, yet some regions contain distinctly human patterns of <u>gene activity</u> that mark the brain's evolution and may contribute to our cognitive abilities, a new Yale-led study has found.

The massive analysis of human, chimpanzee, and monkey tissue published Nov. 23 in the journal *Science* shows that the <u>human brain</u> is not only a larger version of the ancestral primate brain but also one filled with distinct and surprising differences.

"Our brains are three times larger, have many more cells and therefore more processing power than chimpanzee or monkey," said Andre M.M. Sousa, a postdoctoral researcher in the lab of neuroscientist Nenad Sestan and co-lead author of

the study. "Yet there are also distinct small differences between the species in how individual cells function and form connections."

Despite differences in brain size, the researchers found striking similarities between primate species of gene expression in 16 regions of the brain—even in the prefrontal cortex, the seat of higher order learning that most distinguishes humans from other apes. However, the study showed the one area of the brain with the most human-specific gene expression is the striatum, a region most commonly associated with movement.

Distinct differences were also found within regions of the brain, even in the cerebellum, one of the evolutionarily most ancient regions of the brain, and therefore most likely to share similarities across species. Researchers found one gene, ZP2, was active in only human cerebellum—a surprise, said the researchers, because the same gene had been linked to sperm selection by human ova.

"We have no idea what it is doing there," said Ying Zhu, a postdoctoral researcher in Sestan's lab and co-lead author of the paper.

Zhu and Sousa also focused on one gene, TH, which is involved in the production of dopamine, a neurotransmitter crucial to higher-order function and depleted in people living with Parkinson's disease. They found that TH was highly expressed in human neocortex and striatum but absent from the neocortex of chimpanzees.

"The neocortical expression of this gene was most likely lost in a common ancestor and reappeared in the human lineage," Sousa said.

Researchers also found higher levels of expression of the gene MET, which is linked to <u>autism</u> <u>spectrum disorder</u>, in the human <u>prefrontal cortex</u> compared to the other primates tested.



More information: A.M.M. Sousa el al., "Molecular and cellular reorganization of neural circuits in the human lineage," *Science* (2017). <u>science.sciencemag.org/cgi/doi ...</u> <u>1126/science.aan3456</u>

Provided by Yale University APA citation: Small but distinct differences among species mark evolution of human brain (2017, November 23) retrieved 8 June 2022 from <u>https://medicalxpress.com/news/2017-11-small-distinct-differences-species-evolution.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.