

Rods in the retina contribute to daylight vision

November 27 2017

An international research team headed by Thomas Münch from Tübingen University found the contribution of rod photoreceptors in mouse retinas to be much greater than previously assumed. Rods cannot distinguish between colours and were thought to become useless as light levels increase, while vision in daylight conditions is based on cone photoreceptors. The new study – published in *Nature Communications* – shows that rod function can even increase in bright light.

The photoreceptors in the retina, at the back of the eyes, are the primary [light sensitive cells](#) that allow us to see: they convert light into electrical signals. There are two types, rods and cones, and it has generally been assumed that rods are responsible for seeing in dim light conditions, whereas cones mediate [vision](#) in [bright light](#). This division of labour between rods and cones can be found in virtually every biological and medical text book.

A new study challenges this traditional view: A group of researchers from Aalto University, the University of Helsinki and the universities of Tübingen and Manchester shows that [rod photoreceptors](#) do, in fact, contribute to daylight vision. Most surprisingly, their contribution even increases when the daylight becomes brighter, up to the brightest light levels that would ever be encountered in natural environs.

Using transgenic mice without functional cones, the investigators first measured rod-driven signals and could reliably detect them both in the retina and in the brain even at high light levels. Now that they had

isolated these rod-driven signals, they were able to identify them in mice with fully functional cones as well. Their explanation of the mechanism by which rods function in bright light is based on knowledge of light adaptation in rods and the rod visual cycle, the process helping rods to recycle visual pigment after it has been bleached by light.

"We showed that rods can function at bright light," says Thomas Münch, "but it is true that cones can do this much better and much more reliably. However, these new insights may still open new avenues towards treatments for patients without functional cones, so-called rod monochromats." People in modern times are exposed to artificial bright [light](#) for many hours each day. In the old paradigm, this has made reliance on rods for the development of treatments counterintuitive. With these new mechanistic insights into bright rod vision, however, there may be new possibilities for therapies for patients without functional cone vision.

More information: Alexandra Tikidji-Hamburyan et al. Rods progressively escape saturation to drive visual responses in daylight conditions, *Nature Communications* (2017). [DOI: 10.1038/s41467-017-01816-6](#)

Provided by Aalto University

Citation: Rods in the retina contribute to daylight vision (2017, November 27) retrieved 6 May 2023 from <https://medicalxpress.com/news/2017-11-rods-retina-contribute-daylight-vision.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--