

Less frequent biopsy may be option in prostate cancer care

28 November 2017



variable surveillance intervals and competing treatments. There was a delay of three to five months in detecting upgrading with biennial biopsies versus annual biopsies in all cohorts.

"The consequences of more versus less frequent biopsies seem to be similar across cohorts," the authors write.

Several authors disclosed financial ties to the diagnostics industry.

More information: Abstract/Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

(HealthDay)—Biennial biopsies are an acceptable alternative to annual biopsies for men managing low-risk prostate cancer through active surveillance (AS), according to a research published online Nov. 27 in the *Annals of Internal Medicine*.

Lurdes Y.T. Inoue, Ph.D., from University of Washington in Seattle, and colleagues compared the risk for disease progression (upgrading from a Gleason score [GS] of ?6 to ?7) across AS studies after accounting for differences in surveillance protocols in order to evaluate trade-offs of more versus less frequent biopsies. A total of 2,576 men (aged 40 to 80 years) with a GS between 2 and 6 and clinical stage T1 or T2 prostate cancer were included.

The researchers found that estimated risks for biopsy upgrading were similar in the Canary Prostate Active Surveillance Study and University of Toronto studies, but higher in the University of California, San Francisco study and lower in the Johns Hopkins University study, after adjusting for



APA citation: Less frequent biopsy may be option in prostate cancer care (2017, November 28) retrieved 21 November 2022 from https://medicalxpress.com/news/2017-11-frequent-biopsy-option-prostate-cancer.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.