


## Amphetamine abuse disrupts development of mouse prefrontal cortex

8 January 2018



Axons continue to grow to the orbital prefrontal cortex during adolescence. This image illustrates the dual-viral injection method used to label nucleus accumbensprojecting ventral tegmental area neurons. Credit: Hoops et al., eNeuro (2018)

mouse orbital prefrontal cortex is protracted and sensitive to amphetamine in adolescence, eNeuro, DOI: 10.1523/ENEURO.0372-17.2018

More information: Dopamine development in the

Provided by Society for Neuroscience

Recreational drug use during adolescence may disrupt development of an understudied part of the prefrontal cortex, according to a study of male mice published in eNeuro.

Brain cells that utilize the neurotransmitter dopamine are key players in the development of the prefrontal cortex, which continues to mature through early adulthood. During adolescence, the axons of these dopamine neurons extend from the nucleus accumbens to the medial prefrontal frontal cortex.

Cecilia Flores and colleagues show that this delayed growth of dopamine axons also applies to the orbital prefrontal cortex (oPFC), a brain region involved in complex functions such as decisionmaking, but the development of which is poorly understood. Mice exposed to the stimulant amphetamine at doses similar to those obtained when the drug is abused by humans had fewer synapses in the oPFC, likely as a result of reduced axon growth, compared to control mice.



APA citation: Amphetamine abuse disrupts development of mouse prefrontal cortex (2018, January 8) retrieved 28 April 2021 from

https://medicalxpress.com/news/2018-01-amphetamine-abuse-disrupts-mouse-prefrontal.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.