

Prenatal PPI, H2 blocker use linked to asthma risk in child

12 January 2018



increased risk of [childhood asthma](#)," the authors write. "This information may help clinicians and parents to use caution when deciding whether to take acid-suppressing drugs during [pregnancy](#) because of the risk of asthma in offspring."

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Prenatal, maternal, acid-suppressive drug use is associated with an increased risk of childhood asthma, according to a review published online Jan. 11 in *Pediatrics*.

Tianwen Lai, M.D., Ph.D., from Zhejiang University in China, and colleagues conducted a systematic literature review to identify eight population-based studies that assessed acid-suppressive drug use during pregnancy and the risk of childhood asthma in offspring.

In pooled analysis, the researchers found that acid-suppressive drug use in pregnancy was associated with an increased risk of asthma in childhood (relative risk, 1.45). Among [proton pump inhibitor](#) users, the overall risk of asthma in childhood increased (relative risk, 1.34). Similarly, the risk increased in histamine-2 receptor antagonist users (relative risk, 1.57).

"The evidence suggests that prenatal, maternal, acid-suppressive drug use is associated with an

APA citation: Prenatal PPI, H2 blocker use linked to asthma risk in child (2018, January 12) retrieved 10 June 2021 from <https://medicalxpress.com/news/2018-01-prenatal-ppi-h2-blocker-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.