


Gastric bypass and sleeve gastrectomy are comparable for morbid obesity treatment

17 January 2018



Two types of surgery for severe obesity: In the case of gastric bypass, the stomach is bypassed by a part of the small intestine (left), while in the case of the peritoneum, the stomach volume is reduced (right). Credit: USZ, Visceral Surgery

In Switzerland, 5,500 operations to combat morbid obesity are conducted every year. Gastric bypasses and sleeve gastrectomy operations perform similarly: patients lose two-thirds of their excess weight in the long term, as researchers from the University of Basel at the St. Claraspital report in the *Journal of the American Medical Association (JAMA)*. When it comes to gastric acid reflux, the bypass clearly shows better results.

Morbid obesity and associated diseases such as diabetes, cardiovascular diseases and cancer are widespread in the western world. Surgical interventions are currently considered to be the most effective measure in severely obese people with a BMI over 35.

The standard [surgery](#) is a [gastric bypass](#), where part of the stomach and small intestine is bypassed. Other methods are also becoming more popular, such as [sleeve gastrectomy](#), in which the volume of the stomach is reduced. This operation is technically easier to perform and can be done with low complication rate even in patients with

very high BMI.

More than 200 patients examined

To compare these two common types of surgery, the study looked at 217 morbidly obese patients; half were assigned to a gastric bypass and half a sleeve gastrectomy. Their average age was 45.5 years, and 72 percent of them were women. The study was conducted at four locations in Switzerland and was led by Professor Ralph Peterli from the University of Basel at St. Claraspital.

In terms of weight loss, the two types of surgery led to similar results five years after the surgery: the [patients](#) lost 68 percent of their [excess weight](#) after a gastric bypass, and 61 percent after a sleeve gastrectomy. The BMI also fell from 44 to 32 after both operations. Associated diseases such as hypertension, diabetes, altered blood lipids, back/joint pain and depression were also improved, as well as general quality of life.

Improvement in gastric acid reflux

Patients with existing gastric acid reflux benefited more from a gastric bypass operation: after this procedure, symptoms were eliminated in 60 percent, compared to only 25 percent after a sleeve gastrectomy. Sleeve gastrectomy also led to a deterioration or new onset of gastric acid reflux significantly more often. Further surgery or interventions were needed in 22 percent of gastric bypasses and 16 percent of sleeve gastrectomy operations.

More information: Ralph Peterli et al. Effect of Laparoscopic Sleeve Gastrectomy vs Laparoscopic Roux-en-Y Gastric Bypass on Weight Loss in Patients With Morbid Obesity, *JAMA* (2018). [DOI: 10.1001/jama.2017.20897](https://doi.org/10.1001/jama.2017.20897)

Provided by University of Basel

APA citation: Gastric bypass and sleeve gastrectomy are comparable for morbid obesity treatment (2018, January 17) retrieved 25 May 2022 from <https://medicalxpress.com/news/2018-01-gastric-bypass-sleeve-gastrectomy-morbid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.