

Pharmacist participation in stroke response cuts door-to-needle time

19 January 2018



competent pharmacist in the bedside response team for [acute ischemic stroke](#) patients," the authors write.

More information: [Abstract/Full Text](#) ([subscription or payment may be required](#))

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Having a pharmacist at bedside during acute ischemic stroke is associated with significantly shorter door-to-needle (DTN) times, according to a study published recently in the *Annals of Pharmacotherapy*.

Megan A. Rech, Pharm.D., from the Loyola Medical Center in Maywood, Ill., and colleagues conducted a [retrospective cohort study](#) to examine whether [pharmacist](#) presence at bedside during acute ischemic stroke correlated with a reduction in DTN times. Data were included for 125 patients who received recombinant [tissue plasminogen activator](#); 45 and 80 patients had and did not have a pharmacist present (PharmD and no-PharmD groups, respectively).

The researchers found that the median DTN time was significantly shorter in the PharmD group versus the no-PharmD group (48 versus 73 minutes; P

"These findings support the inclusion of a stroke-

APA citation: Pharmacist participation in stroke response cuts door-to-needle time (2018, January 19) retrieved 29 June 2022 from <https://medicalxpress.com/news/2018-01-pharmacist-response-door-to-needle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.