


Scientists describe the mechanism of heart regeneration in the zebrafish

February 12 2018



Section through a zebrafish heart after injury. Cardiomyocytes of the external heart wall are shown in blue. The regenerated tissue visible at the bottom of the image contains some green-stained cardiomyocytes, indicating their origin in the internal part of the heart (red) and their conversion to cells of the myocardial wall. Credit: CNIC

Some animals, including the zebrafish, have a high capacity to regenerate tissues, allowing them to recover fully after cardiac injury. During this process, the heart muscle cells divide to replace the damaged tissue. However, there has been uncertainty about whether all cells contribute equally to the reconstruction of the heart wall. Now, a team of scientists led by Nadia Mercader at the Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC) and the University of Bern (Switzerland), working with collaborators at the University of Zurich (Switzerland), have discovered a high level of plasticity among the cells of the zebrafish heart muscle. The study is published in *Nature Communications*.

After a [heart](#) attack, the human heart loses millions of cardiomyocytes, the cells that form the muscle wall. In contrast, other animal species have a high regenerative capacity, enabling them to replace the injured myocardium with new cardiomyocytes. One such species is the zebrafish (*Danio rerio*). According to first author Héctor Sánchez-Iranzo, the zebrafish "is a widely used model system in cardiovascular research into the mechanisms controlling regeneration, and an inspiration for attempts to develop future regenerative therapies."

The ability of the zebrafish heart to reestablish its function after injury depends on the capacity of its cardiomyocytes to divide and repopulate

the infarcted area, thus eliminating the damaged tissue. Unfortunately, the hearts of most animals, including humans, are unable to activate this process, and therefore after an infarction the [human heart](#) cannot regenerate the lost muscle, which is replaced by non-functional scar tissue.

Cell plasticity

Before the new study, scientists did not know if all cardiomyocytes in the zebrafish heart shared the same regenerative ability or if they were equally able to regenerate all zones of the heart muscle. Cell plasticity, the ability of [cells](#) to convert themselves into another cell type, is frequently observed during embryonic development, but has never before been reported during tissue regeneration in an adult organism.

In the study, the authors investigated two types of cardiomyocyte, one localized in the innermost heart regions, the trabeculae, and the other in the exterior heart wall. Scientists had presumed that during regeneration each cardiomyocyte population would give rise only to the same specialized cell type. But the CNIC study shows that cardiomyocytes from the trabeculae can contribute to the regeneration of the external heart wall. The researchers conclude that their results "reveal a high level of plasticity among [zebrafish](#) cardiomyocytes and that there is more than one way to rebuild a damaged heart."

More information: Héctor Sánchez-Iranzo et al, Tbx5a lineage tracing shows cardiomyocyte plasticity during zebrafish heart regeneration, *Nature Communications* (2018). [DOI: 10.1038/s41467-017-02650-6](https://doi.org/10.1038/s41467-017-02650-6)

Provided by Centro Nacional de Investigaciones Cardiovasculares

Citation: Scientists describe the mechanism of heart regeneration in the zebrafish (2018, February 12) retrieved 6 May 2023 from <https://medicalxpress.com/news/2018-02-scientists-mechanism-heart-regeneration-zebrafish.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.