

Subthreshold phototherapy after birth cuts later phototherapy

February 28 2018



(HealthDay)—Subthreshold phototherapy during birth hospitalization is

associated with reduced readmission for phototherapy, according to a study published online Feb. 26 in *JAMA Pediatrics*.

Andrea C. Wickremasinghe, M.D., from the Kaiser Permanente Santa Clara Medical Center in California, and colleagues estimated the efficacy of subthreshold phototherapy for newborns with total serum bilirubin levels from 0.1 to 3 mg/dL below the appropriate phototherapy threshold during birth hospitalization. Data were included for 25,895 newborns born at 35 or more weeks' gestation.

The researchers found that 19.1 percent of the newborns received subthreshold phototherapy; 4.9 percent of these were readmitted for phototherapy compared with 12.8 percent of the untreated newborns (unadjusted odds ratio, 0.35). The correlation was strengthened after adjustment for confounding variables (odds ratio, 0.28). The estimated numbers needed to treat ranged from 60.8 to 6.3 in the lowest versus the highest quintile of predicted risk. Compared with exclusively breastfed newborns, those who received formula feedings had lower adjusted odds of readmission for phototherapy (odds ratio, 0.58 for >0 to

Citation: Subthreshold phototherapy after birth cuts later phototherapy (2018, February 28) retrieved 10 July 2023 from <https://medicalxpress.com/news/2018-02-subthreshold-phototherapy-birth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.