

Evaluation and treatment of depression may reverse memory and cognitive difficulties

27 March 2018

Individuals with worse depression and mood symptoms are more likely to develop Mild Cognitive Impairment (MCI) and to progress from MCI to dementia. Evaluation and treatment of symptoms of depression may improve or maintain cognitive functioning in some older patients diagnosed with MCI.

MCI is a state in between normal cognition and Alzheimer's disease (AD) [dementia](#), in which the individual has a subjective complaint of memory and other [cognitive](#) difficulties and performance on formal neuropsychological testing is abnormal for age, but these problems do not interfere with routine independent functioning. The majority of people with MCI have progressive memory and cognitive impairment, and eventually are unable to function independently with daily tasks, resulting in a diagnosis of dementia. However, some individuals with MCI do not get progressively worse and some people improve, reverting back from MCI to normal cognition. This study demonstrated that measuring [neuropsychiatric symptoms](#) such as depression, anxiety, agitation and irritability, helps to predict who will progress from normal cognition to MCI, and then who will either progress from MCI to AD dementia or revert back to normal cognition.

In this study, published online in the *Journal of Alzheimer's Disease*, researchers from Boston University School of Medicine (BUSM) analyzed data from 6,763 individuals participating in longitudinal research studies at 34 Alzheimer's Disease Centers across the U.S that are currently or were previously funded by the National Institute on Aging. The data from all centers are entered into a database at the National Alzheimer's Coordinating Center. The participants, whose average age was 72, received annual neurologic examinations and evaluations of their memory and cognitive functioning as well as their neuropsychiatric symptoms. Study partners (e.g., significant others) also rated participants'

neuropsychiatric symptoms and level of functioning. Participants were diagnosed at each evaluation as either cognitively normal, MCI, or AD Dementia by teams of doctors. All participants in this study were cognitively normal at the time of their first examination and were then followed from two to 12 years, with an average of five years.

Results of the study showed that individuals with normal cognition were more likely to progress to MCI if they had more depression, anxiety and other mood symptoms. Similarly, people who had progressed to MCI were more likely to progress even further to AD dementia if they had more of these neuropsychiatric symptoms. An important finding was that approximately one third of the participants who had progressed to MCI reverted back to normal cognition, and that the participants who reverted back had significantly lower neuropsychiatric symptoms and a greater reduction in depression symptoms.

In particular, the researchers noted that improvements in depression and [mood symptoms](#) led to a greater likelihood of cognitive improvement. "The implication is that successfully identifying and providing effective treatment for these neuropsychiatric symptoms, including [depression](#), may potentially improve or maintain [cognitive functioning](#) in many older adults," explained corresponding author Robert Stern, PhD, professor of neurology, neurosurgery and anatomy and neurobiology at BUSM, and Clinical Core Director of the BU Alzheimer's Disease Center. "There are many possible explanations for these findings and further research is needed to address this important issue," Stern cautioned.

Provided by Boston University School of Medicine

APA citation: Evaluation and treatment of depression may reverse memory and cognitive difficulties (2018, March 27) retrieved 25 September 2022 from <https://medicalxpress.com/news/2018-03-treatment-depression-reverse-memory-cognitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.