

Simultaneous chemo and immunotherapy may be better for some with metastatic bladder cancer

11 April 2018

Researchers from Mount Sinai and Sema4, a health information company and Mount Sinai venture, have discovered that giving metastatic bladder cancer patients simultaneous chemotherapy and immunotherapy is safe and that patients whose tumors have certain genetic mutations may respond particularly well to this combination approach, according to the results of a clinical trial published in *European Urology*.

Though chemotherapy and immunotherapy have become standard options for the treatment of metastatic bladder cancer, it was previously unknown whether these therapies could be given together and whether chemotherapy's side effect of weakening the immune system would inhibit immunotherapy. The phase 2 trial was conducted at six cancer centers, and patients in the trial did not show any additional or more severe side effects than patients given chemotherapy or immunotherapy alone, a finding that showed the combination therapy is a safe alternative.

Researchers also generated evidence showing that immunotherapy could boost immune cells in the blood of patients receiving concurrent chemotherapy, allaying previous concerns that chemotherapy might counteract the effects of immunotherapy.

"Because chemotherapy and immunotherapy are the two pillars of treatment for metastatic bladder cancer, we sought to better understand how these treatments might be best given together," said Matthew Galsky, MD, Director of Genitourinary Medical Oncology and Professor of Urology, Medicine, Hematology and Medical Oncology at The Tisch Cancer Institute at the Icahn School of Medicine at Mount Sinai, and Principal Investigator of the study. "Already the results of this trial have inspired the creation of two more trials that seek to

better the treatment of bladder cancer patients by combining chemotherapy with immunotherapy."

One of the new trials, which Dr. Galsky is heading at Mount Sinai and other centers, gives chemotherapy and immunotherapy to a subset of patients with earlier-stage bladder cancer to determine if this combination can head off the need for surgery to remove the bladder, a standard treatment but one with quality-of-life-altering implications that include wearing a urostomy bag outside the body to collect urine. The other trial combines two different chemotherapy regimens with immunotherapy to determine the best types of chemotherapy drugs to combine with immunotherapy.

Dr. Galsky said that the current trial epitomizes the importance of team science. Dr. Galsky, as well as Andrew Uzilov, PhD, Director of Cancer Genomics for Sema4, and Huan Wang, PhD, Sema4 bioinformatics scientist, and other researchers hypothesized that patients with tumors with particular genetic mutations might respond best to the combination of chemotherapy and immunotherapy. And indeed, they found that certain types of mutations in DNA damage response (DDR) genes were associated with better response to the combined chemotherapy and immunotherapy. If validated in subsequent studies, these findings could add a novel biomarker to the "precision oncology toolbox" and refine the selection of patients who might benefit from concurrent administration of chemotherapy and immunotherapy.

"Our work is an example of using genomics to enable precision medicine. By looking for loss-of-function mutations in DDR genes, we may be able to predict who will do well on a combined chemotherapy plus immunotherapy regimen,"



explained Dr. Uzilov. "In our study, we looked at a pool of 55 DDR genes as predictive biomarkers. We are now exploring further which of these genes, and what types of mutations within them, best predict treatment response, as well as the interplay of DDR status with other predictors of immunotherapy response."

Provided by The Mount Sinai Hospital

APA citation: Simultaneous chemo and immunotherapy may be better for some with metastatic bladder cancer (2018, April 11) retrieved 16 June 2022 from

 $\underline{https://medicalxpress.com/news/2018-04\text{-}simultaneous\text{-}chemo\text{-}immunotherapy\text{-}metastatic\text{-}bladder.html}$

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.