

Elder abuse not associated with risk of chronic pain

13 April 2018



between EAN and chronic [pain](#) but indicated cross-sectional associations between the two," the authors write. "This might indicate that, although EAN does not lead to [chronic pain](#), individuals with greater physical limitations are more vulnerable to abuse."

More information: [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Elder abuse and neglect (EAN) is not tied to the risk of chronic pain, according to a study published online March 30 in the *Journal of the American Geriatrics Society*.

Raudah M. Yunus, M.P.H., from University of Malaya in Kuala Lumpur, and colleagues assessed the cross-sectional and longitudinal relationships between EAN and chronic pain in 1,189 community-dwelling, rural Malaysians (aged ≥60 years).

The researchers found that the prevalence of chronic pain was 20.4 percent. Eight factors were significantly associated with chronic pain, including age, education, income, comorbidities, self-rated health, depression, gait speed, and EAN. Abused elderly adults were more likely to have chronic pain (odds ratio, 1.52; 95 percent confidence interval, 1.03 to 2.27), although the significance decreased in longitudinal analyses (risk ratio, 1.14; 95 percent confidence interval, 0.81 to 1.60).

"Our findings indicate no temporal relationship

APA citation: Elder abuse not associated with risk of chronic pain (2018, April 13) retrieved 23 August 2022 from <https://medicalxpress.com/news/2018-04-elder-abuse-chronic-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.