

Slower calorie burn in pregnancy may mean more retained baby weight in obese black moms

April 22 2018



Credit: CC0 Public Domain

Differences in the way women with obesity burn calories during pregnancy may be a contributor to long-term postpartum weight

retention in black moms, according to researchers in Baton Rouge, La. Their new study shows that despite similar levels of food intake and activity levels—and a higher proportion of fat-free mass—obese black women burned fewer calories than their white counterparts. The findings, which suggest a need for more individualized pregnancy weight gain recommendations for obese women, will be presented today at the APS annual meeting at Experimental Biology 2018 in San Diego.

Women with obesity—defined as a body mass index of 30 or above at the start of pregnancy—are at increased risk for gaining too much [weight](#) during their pregnancy and retaining weight after they give birth. In many cases, retained "baby weight" eventually becomes regular weight that may further exacerbate preexisting obesity and related health problems.

Researchers at Louisiana State University's Pennington Biomedical Research Center and the Woman's Hospital in Baton Rouge measured [energy](#) expenditure, body composition and markers of metabolic health in 66 black and white pregnant [women](#) with obesity.

The women were comparable in age (average age 27), BMI (approximately 37) and body fat percentage (approximately 45 percent). The black women were heavier and had a higher amount of fat-free mass (muscle, bone, skin, etc.), which suggest that they had a higher metabolism and would burn more calories. While the black women ate roughly 300 fewer calories per day, the composition of their diet and daily energy expenditure was similar to their white counterparts. However, when the researchers compared energy expenditure—assuming that body mass was the same in black and white women—they found that black women burned roughly 230 fewer calories during the day and 80 fewer calories while sleeping.

The researchers think that the findings may be partly due to the

composition of the fat-free mass in black women. Previous research has shown that while [black women](#) have a higher proportion of calorie-burning muscle, they may have a smaller liver and kidney, two organs that expend more energy than muscle. "Another hypothesis is that African-Americans are more efficient in the use of energy during activity. This can be compared to a more fuel-efficient car, which consumes less gas for the same performance," lead researcher Jasper Most, PhD, of Pennington Biomedical, explained.

Clinicians commonly recommend that pregnant women add an additional 300 to 500 calories per day and that [obese women](#) aim to [gain](#) between 11 and 20 pounds during the course of a normal pregnancy. These findings could have implications on current non-race-specific calorie intake recommendations for pregnant women.

"The [weight gain](#) recommendations for pregnancy are based on weight gain in relation to pregnancy outcomes such as healthy fetal size and in the absence of pregnancy complications. Therefore, our findings cannot immediately contribute to suggestions on healthy weight gain," Most said.

"However, our findings might indeed influence the recommendations for energy intake during pregnancy. The estimated energy-intake requirements to achieve healthy weight gain are the sum of energy that a pregnant woman expends and what she stores (in fat tissue and for fetal growth, for example).

"We show that the [energy expenditure](#) is less in African-American women and therefore energy intake requirements should probably be smaller for African-American women. We estimate if African-American women eat the amount of calories recommended for [pregnant women](#), they would likely gain more weight in pregnancy (up to 7 kg/15 lbs. more) than what is considered healthy," Most said.

Jasper Most, PhD, of Pennington Biomedical, will present the poster "Propensity for excess gestational weight gain in African-American women may be explained by hypometabolic factors in early [pregnancy](#)" on Sunday, April 22, in the Exhibit Hall of the San Diego Convention Center.

Provided by American Physiological Society

Citation: Slower calorie burn in pregnancy may mean more retained baby weight in obese black moms (2018, April 22) retrieved 1 February 2024 from <https://medicalxpress.com/news/2018-04-slower-calorie-pregnancy-retained-baby.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.