


Training compassion 'muscle' may boost brain's resilience to others' suffering

22 May 2018, by Marianne Spoon



As part of a study looking at the ability of compassion meditation to help people cope with the suffering of others, research participants took part in functional MRI brain scans. This allowed scientists to measure activity in regions of the brain involved in dealing with emotional distress. (This photo is not from the study.) PHOTO: David Nevala

It can be distressing to witness the pain of family, friends or even strangers going through a hard time. But what if, just like strengthening a muscle or learning a new hobby, we could train ourselves to be more compassionate and calm in the face of others' suffering?

That is the question behind research from the University of Wisconsin-Madison, and a new study suggests that as little as two weeks of [compassion meditation](#) training—intentionally cultivating positive wishes to understand and relieve the suffering of others—may reduce the distress a person feels when witnessing another's suffering. It may also improve their ability and likelihood to respond with [compassion](#).

The findings, published May 22 in the journal *Frontiers in Psychology*, may have implications for

professions in which people routinely work with others who are suffering, like doctors, law enforcement officers and first responders who experience high levels of distress or empathic burnout.

"Compassion meditation may shift habits of becoming overly distressed when we encounter another's pain," says Helen Weng, assistant professor of psychiatry at the Osher Center for Integrative Medicine at the University of California, San Francisco. She led the research while a graduate student at the Center for Healthy Minds at UW-Madison. "People can learn a calmer and more balanced response when they see someone suffering, even when they are attending more to suffering."

For the study, 24 participants were randomly assigned and trained to do either 30 minutes of compassion meditation or reappraisal training (re-interpreting personally stressful events to decrease negative emotions) every day for two weeks.

The compassion meditation group was trained to visualize people when they were suffering and practice noticing their own personal reactions in a calm and nonjudgmental way. Focusing on a loved one, on themselves, on a stranger, and on someone with whom they had conflict, they also practiced caring for and wishing to help the other person.

In this way, practicing compassion meditation was like exercising a muscle by gradually increasing the "weight" of the relationship with each person considered.

Both groups received brain scans before they were trained and after two weeks of practice to see whether compassion meditation made it easier for those trained to actually look at a suffering person. Humans are visually attentive as a species, Weng says, and looking at someone is a critical first step

in determining if they're in need.

"Your eyes are a window into what you care about," she says. "We wanted to know: Does looking more at suffering in the mind's eye translate into looking more at suffering out in the real world, and can this be done with less distress?"

While in the brain scanner, before and after two weeks of practice, the participants viewed neutral images of strangers as well as emotionally evocative images of people suffering—like a burn victim or crying child. They were instructed to react to the images as they normally would and to make use of their new training.

For instance, people in the compassion training group practiced compassion toward individuals in the images, having thoughts like: "May this person be happy and free from suffering." The reappraisal group reframed the situation: "This person is an actor and isn't really suffering."

The researchers used eye-tracking techniques to record where people spent the most time gazing at each image, whether it was on areas of the image that were more negative—such as the faces of those in suffering—or on less emotionally charged parts of the image. They also compared this to how much time each participant looked at the socially relevant areas of neutral images, like the face of a person walking down the street.

The research team found that the people who had practiced compassion meditation tended to look more directly at suffering in the negative images relative to the neutral photos. They also showed less activity in the amygdala, insula and orbitofrontal cortex—areas of the brain that are usually more active when experiencing emotional distress and might lead to a withdrawal response and averted gaze.

This finding was not present in the reappraisal group, and the results suggest compassion could be a mechanism through which people may become calmer in the face of suffering.

"We communicate a lot with our eyes, and this research suggests that compassion training has an

impact on the body and can actually shift where you direct your visual attention when you see others in pain," says Weng.

Compassion meditation slows things down so people can practice being more calm, notice the feelings that arise and learn to be less reactive, she adds. "This gives you more mental space to focus on the other person, to practice wishing kindness and wanting them to be well, and I think both parts are really important for effectively responding to people suffering."

Though the results are exciting, Weng says a larger and more diverse sample of people should be involved in a repeated study. The study was also performed with people who had never meditated before and it's not yet clear whether scientists would find more pronounced results in people who already had extensive compassion meditation practice.

Compassion [meditation](#) could also be used as a strategy for working with [people](#) with conditions that affect how comfortable they are making eye contact with others, says Richard Davidson, founder of the Center for Healthy Minds and senior author of the study.

"The pattern of these findings—an increase in looking at suffering while simultaneously down-regulating neural circuits associated with negative emotion—is a winning combination that may be beneficial for a wide range of conditions including autism and social anxiety disorder in which gaze aversion and social discomfort are hallmark signs," he adds.

More information: Helen Y. Weng et al. Visual Attention to Suffering After Compassion Training Is Associated With Decreased Amygdala Responses, *Frontiers in Psychology* (2018). [DOI: 10.3389/fpsyg.2018.00771](https://doi.org/10.3389/fpsyg.2018.00771)

Provided by University of Wisconsin-Madison

APA citation: Training compassion 'muscle' may boost brain's resilience to others' suffering (2018, May 22) retrieved 1 December 2022 from <https://medicalxpress.com/news/2018-05-compassion-muscle-boost-brain-resilience.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.