

Link between autoimmune disorders and psychosis confirmed in new study

29 June 2018, by Alexis E Cullen



Lid retraction is a typical symptoms of Graves' disease.
Credit: [Jonathan Trobe/Wikimedia Commons, CC BY](#)

People with autoimmune disorders, a collection of diseases where the body's immune system attacks its own cells, are more likely to have psychosis, according to our [latest research](#).

Previous research found that rates of [rheumatoid arthritis](#) were lower in people with [psychosis](#) than would be expected in the general population. But later studies showed that other [autoimmune disorders](#), such as [coeliac disease](#) and [autoimmune thyroid disorders](#), were more common in people with psychosis. This led scientists to the view that there is a connection between autoimmune [disorders](#) and psychosis. But conflicting findings meant that it was difficult to reach any conclusions about the [relationship](#).

Given the uncertainty about the relationship between these disorders and psychosis, and growing interest in this area, we decided to review the research and conduct a meta-analysis – a method in which data from several studies are combined and analysed together to give a more statistically robust result than individual studies can

provide.

Our study, which is [published in *Biological Psychiatry*](#), included 30 relevant studies, and contained data on 25m people.

What we found

We focused on autoimmune disorders that affect the peripheral system, such as type 1 diabetes, as we were particularly interested in whether autoimmune disorders that target the body, as opposed to the brain, could still influence the development of psychosis.

For our main analysis, we combined data from all non-neurological autoimmune disorders, except for rheumatoid [arthritis](#) (given the well-established negative association reported with psychosis) and found that, overall, people with any autoimmune disorder were 40% more likely to have a psychotic disorder, such as schizophrenia.

For our secondary analysis, we examined individual autoimmune disorders. We found the likelihood of having psychosis was higher for [pernicious anaemia](#), [pemphigoid](#) (a disease characterised by skin blisters), [psoriasis](#), coeliac disease and [Graves' disease](#) (the disease suffered by [Marty Feldman](#) that causes protrusion of the eyeballs). But it was lower for [rheumatoid arthritis](#) and [ankylosing spondylitis](#) (a type of arthritis that mainly affects the spine), suggesting that these disorders are protective.

Looking for causes

There are a range of possible mechanisms that might underlie the relationships we found. Given that people with psychosis have also been found to show higher levels of [inflammatory markers](#) in the blood than healthy people, and that inflammation is a core feature of autoimmune disorders, inflammation is a likely candidate.

But rheumatoid arthritis and ankylosing spondylitis are also characterised by higher levels of inflammation, so this would not explain the negative relationships we found with these disorders.

Although all autoimmune disorders activate the body's immune system, the exact response differs depending on the disorder. This might go some way to explaining why we found different relationships for individual autoimmune disorders, and suggests that inflammation cannot be the only mechanism.

because early intervention has been [shown to improve long-term outcomes](#) for people in the initial stages of a psychotic disorder.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

It is possible that there might be a genetic link between autoimmune disorders and psychosis. In fact, [research has recently shown](#) that variations within specific genes are associated with both schizophrenia and rheumatoid arthritis. That is, people with one variation of the gene are at risk for schizophrenia, while people with the other variation are at risk for rheumatoid arthritis. This might explain why rheumatoid arthritis appears to be protective for psychosis.

Newly discovered antibodies (part of the immune system's armoury) that go rogue and mistakenly attack brain cells [might also explain the link](#). These sorts of antibodies are thought to cause psychotic symptoms, such as paranoia and hallucinations, in some people.

Although our study cannot tell us why autoimmune disorders and psychosis occur together more commonly than we would expect, it provides stronger evidence that a relationship does exist.

Our group is involved in further research to help us better understand the mechanisms that might underlie this complex relationship.

Early intervention

So what is the real-world application of these findings? Although the risk of psychosis is only slightly increased for people with autoimmune disorders, our findings suggest that perhaps doctors ought to monitor people with certain autoimmune disorders for early signs of psychosis – especially pernicious anaemia, Graves' disease and pemphigoid, which showed the most consistent relationships with psychosis. This is important

APA citation: Link between autoimmune disorders and psychosis confirmed in new study (2018, June 29) retrieved 11 June 2022 from <https://medicalxpress.com/news/2018-06-link-autoimmune-disorders-psychosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.