


# Novel PET imaging method more fully evaluates extent of rheumatoid arthritis inflammation

9 July 2018



CT and  $^{11}\text{C}$ -PBR28 PET SUV images of both knees of (i) an RA patient with clinical signs of synovitis both knees, and (ii) a healthy control. Credit: N Narayan et al. 2018, University of Oxford, Headington, Oxford, UK

A new positron emission tomography (PET) imaging method more fully evaluates the extent of rheumatoid arthritis by targeting translocator protein (TSPO) expression in the synovium (joint lining tissue). The study is featured in the July issue of *The Journal of Nuclear Medicine*.

PET radioligands that target TSPO, which is highly expressed on activated [macrophages](#) (immune cells responding to inflammation), have proved an excellent tool for imaging joint inflammation in rheumatoid arthritis (RA).

"TSPO-targeted imaging has long been used as a means of imaging macrophage infiltration in vivo," explains Nehal Narayan of the University of Oxford in Oxford, U.K. "Numerous studies have demonstrated TSPO-targeted PET as a highly sensitive and specific means of imaging synovitis [inflammation of joint lining tissue], purportedly through imaging synovial macrophage infiltration, a critical process in RA pathogenesis. However, this premise does not take into account the ubiquitous expression of TSPO."

She points out, "Here we present the first ever analysis of TSPO expression in the major constituents of RA pannus (inflamed synovium), demonstrating that TSPO PET likely acts as an imaging tool of not only macrophages, but also activated synovial fibroblasts, a cell group increasingly recognised to play a critical role in RA inflammation."

The study included three RA patients and three healthy volunteers who underwent PET scans of both knees using the TSPO radioligand carbon-11 ( $^{11}\text{C}$ )-PBR28. In addition, cellular expression of TSPO was examined in synovial tissue from these individuals, plus three more RA patients and three more healthy patients (undergoing knee arthroscopy for injuries). TSPO mRNA expression and hydrogen-3 ( $^3\text{H}$ )-PBR28 radioligand binding were assessed using in vitro monocytes, macrophages, fibroblast-like synoviocytes (FLS) and CD4+ T-lymphocytes.

Results showed that the  $^{11}\text{C}$ -PBR28 PET signal was significantly higher in RA joints compared to healthy joints. In addition,  $^3\text{H}$ -PBR28 specific binding in synovial tissue was approximately 10-fold higher in RA patients compared to healthy controls. Immunofluorescence revealed TSPO expression on macrophages, FLS and CD4+ T cells. In vitro study demonstrated highest TSPO mRNA expression and  $^3\text{H}$ -PBR28 specific binding in activated FLS, non-activated and activated 'M2' reparative macrophages. The lowest TSPO expression was in activated and non-activated CD4+ T lymphocytes.

Narayan notes, "It is well recognised that not all currently available treatments are capable of controlling joint inflammation in all patients with [rheumatoid arthritis](#), hence the need to develop new pharmacological therapies. This work

demonstrates that TSPO PET is able to act as a means of imaging not only synovial macrophages, but also activated synovial fibroblasts. The crucial role of the fibroblast and its soluble products in RA pathogenesis is increasingly realised."

She adds, "Indeed, there has been recent interest in targeting activated fibroblasts as a novel targeted treatment strategy for RA. Therefore, TSPO PET imaging in early phase clinical trials may provide a sensitive indication of treatment response to such novel therapies with a view to informing the design of later stage clinical trials. As our knowledge of cellular TSPO [expression](#) and behavior grows, TSPO-targeted imaging may also give us unique insights into the pathogenesis of inflammatory disease."

**More information:** Nehal Narayan et al, Translocator Protein as an Imaging Marker of Macrophage and Stromal Activation in Rheumatoid Arthritis Pannus, *Journal of Nuclear Medicine* (2018). [DOI: 10.2967/jnumed.117.202200](https://doi.org/10.2967/jnumed.117.202200)

Provided by Society of Nuclear Medicine and Molecular Imaging

APA citation: Novel PET imaging method more fully evaluates extent of rheumatoid arthritis inflammation (2018, July 9) retrieved 28 April 2021 from <https://medicalxpress.com/news/2018-07-pet-imaging-method-fully-extent.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*