


Cetuximab+RT found to be inferior to standard treatment in HPV+ oropharyngeal cancer

August 14 2018

Parts of the Oropharynx



© 2016 Terese Winslow LLC
U.S. Govt. has certain rights

Anatomical illustration of the oropharynx. Credit: National Cancer Institute

An interim analysis of data from a randomized clinical trial of patients with human papillomavirus (HPV)-positive oropharyngeal cancer found that treatment with radiation therapy and cetuximab is associated with worse overall and progression-free survival compared to the current standard treatment with radiation and cisplatin. The trial was designed to see if cetuximab with radiation would be less toxic than cisplatin with radiation without compromising survival for patients with the disease.

The phase 3 trial, which closed enrollment in 2015, was funded by the National Cancer Institute (NCI), part of the National Institutes of Health, and led by NRG Oncology, part of NCI's National Clinical Trials Network. The data monitoring committee overseeing the trial recommended releasing the data after an interim data analysis showed that [cetuximab](#) with radiation was associated with inferior overall and progression-free survival, compared to [cisplatin](#) and radiation. The U.S. Food and Drug Administration has previously approved cetuximab with radiation for patients with head and neck cancer, including oropharyngeal cancer. Cetuximab with radiation is an accepted standard of care, especially for patients who cannot tolerate cisplatin.

"The goal of this trial was to find an alternative to cisplatin that would be as effective at controlling the cancer, but with fewer side effects," said Andy Trotti, M.D., of the Moffitt Cancer Center in Tampa, Florida, a lead investigator of the trial. "We were surprised by the loss of tumor control with cetuximab."

There has been a lot of recent interest in the cancer clinical research community in evaluating the "de-escalation" of therapies for cancers that have a good prognosis, such as HPV-positive cancer of the oropharynx (the part of the throat at the back of the mouth, including the soft palate, the base of the tongue, and the tonsils). The goal is to improve patients' quality of life and reduce long-term toxic effects without compromising treatment efficacy. HPV-positive oropharyngeal cancer is frequently

diagnosed in individuals in their 50s and 60s, and is associated with high survival rates, providing the incentive for this trial. Moreover, the incidence of this type of cancer has increased rapidly in recent years in the United States.

"Clinical [trials](#) designed to test less toxic treatment strategies for patients without compromising clinical benefit are a very important area of interest for NCI and the cancer research community," said Shakun Malik, M.D., of NCI's Division of Cancer Treatment and Diagnosis.

This trial's primary objective was to determine whether the substitution of cetuximab for cisplatin with radiation would result in comparable overall survival while reducing toxic side effects with improved long-term quality of life. The trial enrolled 849 patients with HPV-positive oropharyngeal cancer who were randomly assigned to receive either cetuximab or cisplatin with radiation. The study had three planned interim analyses.

The third and final interim analysis, done after a median follow-up of 4.5 years, found that overall survival on the cetuximab arm was significantly inferior to the cisplatin arm. Overall rates of serious (grade 3-5) adverse events were similar for patients in both groups. However, as the researchers expected, toxic side effects were different, with adverse events of renal toxicity, hearing loss, and bone marrow suppression more common in patients in the cisplatin arm, while body rash was more common in the cetuximab arm. All patients in this trial had completed therapy at the time of this analysis.

"This trial is the first [randomized clinical trial](#) specifically designed for [patients](#) with HPV-positive oropharyngeal [cancer](#), and it establishes cisplatin with [radiation](#) as the standard of care," said Maura Gillison, M.D, Ph.D., of the University of Texas MD Anderson Cancer Center in Houston, the other lead investigator of the trial.

More information: clinicaltrials.gov/ct2/show/NCT01302834

Provided by National Cancer Institute

Citation: Cetuximab+RT found to be inferior to standard treatment in HPV+ oropharyngeal cancer (2018, August 14) retrieved 25 February 2023 from <https://medicalxpress.com/news/2018-08-cetuximabrt-inferior-standard-treatment-hpv.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.