

People with fibromyalgia have inflammation of the brain

26 September 2018, by Felicia Lindberg



Credit: CC0 Public Domain

The causes of the difficult-to-treat pain syndrome fibromyalgia are largely unknown. Using PET brain imaging, researchers at Karolinska Institutet and Massachusetts General Hospital have now shown that glial cells – the central nervous system's immune cells – are activated in the brains of patients with fibromyalgia. The finding has been published in the scientific journal *Brain, Behavior, and Immunity* and may open the way for new therapies.

Fibromyalgia is a chronic [pain](#) syndrome that causes extensive pain in the muscles and joints, severe fatigue, insomnia and cognitive difficulties. The higher pain sensitivity that is characteristic of the syndrome has been related to functional and structural alterations of [brain](#) regions associated with pain processing.

In 2012, Eva Kosek's research group at Karolinska Institutet showed that patients with fibromyalgia had elevated levels of certain inflammatory substances (cytokines) in the cerebrospinal fluid, suggesting inflammation of the central nervous system. Their findings were subsequently corroborated by other researchers, but the source

of the inflammation remained unknown.

Glial cells are activated

Using modern PET (positron-emission topography) brain imaging Eva Kosek's team has now been able to show that the central nervous system's immune [cells](#), called [glial cells](#), are activated and thus give rise to inflammation of the brain.

"As far as we know, this is the first time it's been shown that glial cells are involved in the pathogenesis of fibromyalgia," says Professor Eva Kosek from the Department of Clinical Neuroscience, Karolinska Institutet.

The results show that in Swedish and American patients with fibromyalgia, glial cells are activated in large parts of the cerebral cortex, and that the degree of activation was related to the degree of fatigue reported by the patients.

Objective aberrations in the brain

"The findings may open the way for the development of completely new therapies for this currently difficult-to-treat condition," says Professor Kosek. "The fact that scientific research is able to demonstrate objective aberrations in the brains of people with fibromyalgia will hopefully mitigate the suspicion with which patients are often treated by the health services and society."

Today, an estimated 200,000 Swedes, mainly women, suffer from [fibromyalgia](#). The brains of people with the condition are known to have an impaired ability to dampen pain signals, which means that things that are normally painless cause considerable discomfort.

The study was a collaboration between Eva Kosek's research group at Karolinska Institutet in Sweden, the PET Centre at Karolinska Institutet and Dr. Marco Loggia's research group at Martinos

Center for Biomedical Imaging at Massachusetts General Hospital, USA. The research in Sweden was financed from several sources, including the EU's 7th Framework and Programme and a donation from the Lundblad Family. The Swedish part of the project was also funded by Stockholm County Council, the Swedish Research Council, the Swedish Rheumatism Association and the Swedish Fibromyalgia Association.

More information: Daniel S. Albrecht et al. Brain glial activation in fibromyalgia – A multi-site positron emission tomography investigation, *Brain, Behavior, and Immunity* (2018). DOI: [10.1016/j.bbi.2018.09.018](https://doi.org/10.1016/j.bbi.2018.09.018)

Provided by Karolinska Institutet

APA citation: People with fibromyalgia have inflammation of the brain (2018, September 26) retrieved 25 May 2022 from <https://medicalxpress.com/news/2018-09-people-fibromyalgia-inflammation-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.