


Ketamine is a safe, effective alternative to opioids in treating acute pain in the ED

15 October 2018



Systematic review and meta-analysis: Randomized controlled trials 1946-2017; adult emergency department patients. Credit: KIRSTY CHALLEN, B.SC., MBCHB, MRES, PH.D., LANCASHIRE TEACHING HOSPITALS, UNITED KINGDOM.

Intravenous, low-dose ketamine (LDK) is as effective as intravenous morphine in the control of acute pain in adults in the emergency department (ED). That is the finding of a study to be published in the October 2018 issue of *Academic Emergency Medicine* (AEM), a journal of the Society for Academic Emergency Medicine (SAEM). The results indicate that ketamine can be considered as an alternative to opioids for ED short-term pain control.

The lead author of the study is Nicholas Karlow, MPHS, a medical student at the Washington University School of Medicine in St. Louis, Missouri. The findings of the study are discussed in the featured [episode](#) of SGEM Hop (Skeptics Guide to EM Hot Off the Press).

The systematic review and meta-analysis by

Karlow, et al. maintains that there is a role for opioids in the treatment of pain in the ED, but suggest that as physicians continue to face pressure to reduce opioid use, it is important to establish that alternatives such as ketamine are comparable in providing patients with appropriate analgesia in a similar time frame.

The study further suggests that for patients with opioid use disorders or substance use disorders that require a potent analgesic in the [emergency department](#), ketamine may be a favorable option compared to an opioid.

Moving forward, the authors suggest that observational studies assessing adverse events should use similar outcome measures and time frames, and that researchers should explore patient and physician satisfaction with ketamine analgesia and side effects compared to other opioid alternatives for [acute pain](#).

"Karlow and colleagues provide persuasive evidence that emergency physicians can reasonably expect sub-dissociative ketamine to be as effective as morphine for patients with acute abdominal or musculoskeletal pain. Minor ketamine adverse effects will likely prevent this therapy from becoming routinely first line, but low dose ketamine represents a good alternative choice for selected patients," commented Steven M. Green, MD, professor of [emergency](#) medicine and residency director at Loma Linda University, California.

Dr. Green's principal research interest has been on procedural sedation and analgesia, with numerous studies of [ketamine](#) dating back to 1990 and more recent works relating to sedation's optimal practice, politics, and future. He is a deputy editor at *Annals of Emergency Medicine* journal.

More information: Nicholas Karlow et al, A Systematic Review and Meta-analysis of Ketamine as an Alternative to Opioids for Acute Pain in the

Emergency Department, *Academic Emergency
Medicine* (2018). DOI: [10.1111/acem.13502](https://doi.org/10.1111/acem.13502)

Provided by Society for Academic Emergency
Medicine

APA citation: Ketamine is a safe, effective alternative to opioids in treating acute pain in the ED (2018, October 15) retrieved 26 April 2021 from <https://medicalxpress.com/news/2018-10-ketamine-safe-effective-alternative-opioids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.