

## Aspirin, anticoagulants similarly prevent VTE after TKA

19 October 2018



those who received no pharmacologic prophylaxis, 1.16 percent treated with aspirin alone, 1.42 percent of those treated with anticoagulation alone, and 1.31 percent prescribed both aspirin and anticoagulation. For the composite of VTE or death, aspirin only was non-inferior versus other chemoprophylaxis (adjusted odds ratio, 0.85; 95 percent confidence interval [CI], 0.68 to 1.07; P for inferiority = 0.007). Bleeding occurred in 1.1 percent of patients overall and specifically in 1.5 percent without prophylaxis, 0.9 percent in the aspirin group, 1.14 percent with anticoagulation, and 1.35 percent of those receiving both. For bleeding complications, aspirin alone was also noninferior (adjusted odds ratio, 0.8; 95 percent CI, 0.63 to 1; P for inferiority

"In this study of patients undergoing TKA, aspirin was not inferior to other anticoagulants in the postoperative rate of VTE or death," the authors write.

More information: Abstract/Full Text (subscription or payment may be required)
Editorial (subscription or payment may be required)

Copyright © 2018 HealthDay. All rights reserved.

(HealthDay)—Among patients undergoing total knee arthroplasty (TKA), aspirin alone may provide protection against postoperative venous thromboembolism (VTE) that is similar to that of other anticoagulants, according to a study published online Oct. 17 in *JAMA Surgery*.

Brandon R. Hood, M.D., from the University of Michigan in Ann Arbor, and colleagues retrospectively assessed the acceptability of aspirin alone versus anticoagulant prophylaxis for reducing the risk of postoperative VTE among 41,537 patients who underwent primary TKA between April 1, 2013, and Oct. 31, 2015. Pharmacologic prophylaxis methods were: neither aspirin nor anticoagulants (668 patients), aspirin only (12,831 patients), anticoagulant only (e.g., low-molecular-weight heparin, warfarin, and Xa inhibitors; 22,620 patients), and both aspirin and anticoagulant (5,418 patients).

The researchers found that VTE events occurred in 1.38 percent of patients, including 4.79 percent of



APA citation: Aspirin, anticoagulants similarly prevent VTE after TKA (2018, October 19) retrieved 25 August 2022 from <a href="https://medicalxpress.com/news/2018-10-aspirin-anticoagulants-similarly-vte-tka.html">https://medicalxpress.com/news/2018-10-aspirin-anticoagulants-similarly-vte-tka.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.