

Psoriasis linked to increased risk for inflammatory bowel disease

30 October 2018



symptoms," conclude the authors.

More information: Abstract/Full Text

Copyright © 2018 HealthDay. All rights reserved.

(HealthDay)—Psoriasis appears to be significantly associated with inflammatory bowel disease (IBD), according to a review published online Oct. 24 in *JAMA Dermatology*.

Yun Fu, M.D., from Chang Gung Memorial Hospital in Taiwan, and colleagues assessed case-control, cross-sectional, or cohort studies that examined either the odds or risk for IBD in patients with psoriasis.

Based on five case-control or cross-sectional studies and four cohort studies that involved a total of 7,794,087 participants, the researchers found significant associations between psoriasis and Crohn's disease (odds ratio, 1.7) and between psoriasis and ulcerative colitis (OR, 1.75). In addition, patients with psoriasis had an increased risk for Crohn's disease (risk ratio, 2.53) and ulcerative colitis (risk ratio, 1.71).

"Gastroenterology consultation may be indicated when patients with <u>psoriasis</u> present with bowel



APA citation: Psoriasis linked to increased risk for inflammatory bowel disease (2018, October 30) retrieved 21 September 2022 from https://medicalxpress.com/news/2018-10-psoriasis-linked-inflammatory-bowel-disease.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.