

Research: A third of hospitalised adolescents with life-threatening anorexia are not thin

December 3 2018



Credit: CC0 Public Domain

Health workers are being urged to closely monitor adolescents losing weight after a study of patients with anorexia nervosa found 31 per cent had all the cognitive features and physical complications of the disease

without being underweight.

Dietitian Melissa Whitelaw is calling for a change to [anorexia](#) nervosa's diagnostic criteria after finding that patients with "atypical anorexia" suffer serious health concerns despite being within or above the healthy [weight](#) range.

Her study, published in the *Journal of Adolescent Health*, looked at 171 patients aged 12-19 admitted to the Royal Children's Hospital's eating disorder program between 2005-2013 suffering anorexia nervosa. She found:

- 51 of the patients were "atypical" with significant eating disorder psychopathology, but not [underweight](#)
- Rather than being underweight, greater [weight loss](#) was associated with life threateningly low pulse rates, a complication of starvation in anorexia nervosa that requires admission
- Those with atypical anorexia nervosa also suffered [low blood pressure](#) and deranged blood electrolytes
- Importantly, no complication was independently associated with underweight, the hallmark of anorexia
- No participant in the study was being monitored by a health professional for weight loss, their relationship with food, or their methods of losing weight.

Mrs Whitelaw said atypical patients may have been encouraged by family or [health professionals](#) to lose weight, which frequently resulted in positive re-enforcement and encouragement about how good they looked, praise for losing weight and the ability to wear trendier clothes, which spurred them on to try to lose more weight.

Atypical anorexia nervosa patients might have lost about a quarter of their body weight, but the body could go into "starvation mode" if as

little as 10 per cent of weight was lost quickly, causing the heart rate to slow to preserve energy.

"If adolescents lose weight, it doesn't matter what weight they are, a health professional should monitor them to check that weight loss is appropriate and if so, that it is done gradually. They should also monitor the adolescent's dietary intake and relationship with food and exercise for signs the patient was spiralling into an [eating disorder](#). Following large amounts of weight loss, careful medical assessment is also recommended," Mrs Whitelaw said.

Once a person entered starvation mode the only way to increase the heart rate was re-feeding and weight gain, which in this cohort, required hospitalisation.

Mrs Whitelaw said people could understand an extremely thin patient needing to gain weight, but it was often a shock to individuals and families when someone within or above the healthy weight range needed to gain weight.

Mrs Whitelaw said atypical anorexia nervosa was commonly perceived as less severe than anorexia nervosa, but her research showed the health consequences could be just as dangerous and it was time to change the current [diagnostic criteria](#) which stated those with anorexia nervosa must be underweight.

"What we are seeing now is that you can have a healthy body weight but be just as sick as someone with typical anorexia nervosa, including having the same thoughts about eating and food," she said. "We need to redefine anorexia because an increasing proportion of [anorexia nervosa](#) patients are atypical and more difficult to recognise. The definition should refer to weight loss, not just underweight."

Mrs Whitelaw said: "The face of eating disorders is changing against a backdrop of increasing prevalence of overweight and obesity. Families, teachers, sports coaches and others interacting with young people should not delay seeking help for adolescents with worrying eating patterns if they have lost weight, even if they are not underweight."

More information: Melissa Whitelaw et al. Predictors of Complications in Anorexia Nervosa and Atypical Anorexia Nervosa: Degree of Underweight or Extent and Recency of Weight Loss?, *Journal of Adolescent Health* (2018). [DOI: 10.1016/j.jadohealth.2018.08.019](https://doi.org/10.1016/j.jadohealth.2018.08.019)

Provided by University of Melbourne

Citation: Research: A third of hospitalised adolescents with life-threatening anorexia are not thin (2018, December 3) retrieved 16 March 2023 from <https://medicalxpress.com/news/2018-12-hospitalised-adolescents-life-threatening-anorexia-thin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.