

Risk factors for falling are different in older men versus older women

5 December 2018



different fall risk profiles. If we take these <u>sex</u> differences into consideration in screening and prevention programs for falls, we may potentially enable a better person-centered approach."

More information: Stina Ek et al, Risk Factors for Injurious Falls in Older Adults: The Role of Sex and Length of Follow-Up, *Journal of the American Geriatrics Society* (2018). DOI: 10.1111/jgs.15657

Provided by Wiley

Credit: CC0 Public Domain

Different factors were associated with falling in men versus women in a study of 3,112 communitydwelling adults aged 60 years and older. The findings are published in the *Journal of the American Geriatrics Society*.

In the short term, living alone, needing help with activities of daily living, and previous falls were independently associated with injurious falls in women. Low systolic blood pressure, impaired ability to rise from a chair, and previous falls were associated with injurious falls in men.

Long-term risk factors were underweight, <u>cognitive</u> <u>impairment</u>, use of medications associated with an increased risk of falling, and needing help with activities of daily living for women and smoking, heart disease, impaired balance, and a previous fall for men.

"Despite the numerous studies on <u>risk factors</u> for falls in <u>older adults</u>, studies examining sex differences are scarce," said lead author Stina Ek, of the Karolinska Institutet, in Sweden. "The results of our study suggest that men and women have



APA citation: Risk factors for falling are different in older men versus older women (2018, December 5) retrieved 11 August 2022 from https://medicalxpress.com/news/2018-12-factors-falling-older-men-women.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.